

U.S. Fish & Wildlife Service

Abundance and Run Timing of Adult Salmon in the Gisasa River, Koyukuk National Wildlife Refuge, Alaska, 2022

Alaska Fisheries Data Series Number 2024–2



**Fairbanks Fish and Wildlife Conservation Office
Fairbanks, Alaska
April 2024**



The Alaska Region Fisheries Program of the U.S. Fish and Wildlife Service conducts fisheries monitoring and population assessment studies throughout many areas of Alaska. Dedicated professional staff located in Anchorage, Fairbanks, and Kenai Fish and Wildlife Offices and the Anchorage Conservation Genetics Laboratory serve as the core of the Program's fisheries management study efforts. Administrative and technical support is provided by staff in the Anchorage Regional Office. Our program works closely with the Alaska Department of Fish and Game and other partners to conserve and restore Alaska's fish populations and aquatic habitats. Our fisheries studies occur throughout the 16 National Wildlife Refuges in Alaska as well as off-Refuges to address issues of interjurisdictional fisheries and aquatic habitat conservation. Additional information about the Fisheries Program and work conducted by our field offices can be obtained at: <https://www.fws.gov/about/region/alaska>

The Alaska Region Fisheries Program reports its study findings through the Alaska Fisheries Data Series (AFDS) or in recognized peer-reviewed journals. The AFDS was established to provide timely dissemination of data to fishery managers and other technically oriented professionals, for inclusion into agency databases, and to archive detailed study designs and results for the benefit of future investigations. Publication in the AFDS does not preclude further reporting of study results through recognized peer-reviewed journals.

Cover Photo: Gisasa River Weir, 2022. Photo courtesy of Nicole Farnham.

Disclaimer: The findings and conclusions in this article are those of the author(s) and do not necessarily represent the views of the U.S. Fish and Wildlife Service. The use of trade names of commercial products in this report does not constitute endorsement or recommendation for use by the federal government.

Abundance and Run Timing of Adult Salmon in the Gisasa River, Koyukuk National Wildlife Refuge, Alaska, 2022

Nicole Farnham, Jeremy Carlson, and Naomi Brodersen

Abstract

A resistance board weir was operated by the U.S. Fish and Wildlife Service, Northern Alaska Fish and Wildlife Field Office to collect information on abundance, run timing, and biology of returning adult Chinook Salmon *Oncorhynchus tshawytscha* and summer Chum Salmon *O. keta* in the Gisasa River. The weir has been operated at this location since 1994 except for 2018, 2020, and 2021. This is the fifth year video technology has been incorporated into the project since 2015. In 2022, the weir was operated from the end of the day on June 29 through the morning of July 29 during which an estimated 503 Chinook Salmon and 3,300 summer Chum Salmon passed through the weir. Other species enumerated through the weir, in order of abundance, were Northern Pike *Esox lucius* ($N = 22$), Longnose Sucker *Catostomus catostomus* ($N = 10$), Broad Whitefish *Coregonus nasus* ($N = 9$), Sockeye Salmon *O. nerka* ($N = 5$), Humpback Whitefish *Coregonus pidschian* ($N = 3$), Arctic Grayling *Thymallus arcticus* ($N = 1$), and Pink Salmon *O. gorbuscha* ($N = 1$). A total of 42 Chinook Salmon and 294 summer Chum Salmon were sampled for age, sex, and length (ASL) data. The sample size goal was not met for Chinook Salmon and ASL analyses could not be performed for this species. The summer Chum Salmon run was composed of four strata (July 4 through 13, July 14 through 17, July 18 through 24, and July 25 through 28). The percent female for all strata combined was 55. There were two primary age classes identified for summer Chum Salmon, 0.3 and 0.4. Length-at-age of male summer Chum Salmon was larger than females.

Introduction

The Gisasa River, located within the Koyukuk National Wildlife Refuge in northcentral interior Alaska, is a tributary of the Koyukuk River and provides spawning and rearing habitat for Chinook Salmon *Oncorhynchus tshawytscha* and summer Chum Salmon *O. keta*. These salmon species in the Gisasa River contribute to mixed stock subsistence and commercial fisheries in the Yukon River (USFWS 1993). The U.S. Fish and Wildlife Service (USFWS), through Section 302 of the Alaska National Interest Lands Conservation Act, has a responsibility to ensure that salmon populations within federal conservation units are conserved in their natural diversity, international treaty agreements are met, and subsistence opportunities are maintained.

Yukon River salmon returns declined in the late 1990s (Kruse 1998). These declines led to harvest restrictions, complete fishery closures, and spawning escapements below management goals (Vania et al. 2002). Since the late 1990s, summer Chum Salmon returns have shown considerable variability with no pattern of declining numbers, although 2021 was the lowest summer Chum Salmon return on record (JTC 2022). However, Chinook Salmon returns rebounded and continued to improve from 2001 to 2006, and then declined again from 2007 to 2016. The 2017 passage was the highest since 2003, while the 2021 passage was the lowest Chinook Salmon return since 2003 (JTC 2022). In the Yukon River drainage, management of individual stocks does not occur, and accurate escapement

data are limited. In-season management of the salmon fisheries is conducted using preseason projections based on parent stock returns; salmon counts from sonar projects at Pilot Station and Eagle; data from test fisheries and escapement assessment projects; and harvest data from subsistence and commercial fisheries.

Historically, escapement information on individual salmon stocks from the Koyukuk River has been collected by aerial surveys. The Alaska Department of Fish and Game (ADF&G) has conducted these surveys on several index tributaries within the Koyukuk River drainage intermittently since 1960 (Barton 1984). Aerial survey results, however, are highly variable and provide only a point in time index of relative run strength. Counts produced using weirs or counting towers provide a better estimation of escapement, and weirs provide a platform for collecting other biological data. Weirs or counting towers have been operated on five different Koyukuk River tributaries since 1994, and weirs currently operate on the Gisasa River and at Henshaw Creek (Figure 1).

The USFWS, Northern Alaska Fish and Wildlife Field Office (NAFWFO), has operated a resistance board weir on the Gisasa River since 1994 (Melegari and Wiswar 1995) (Appendix 1). Chinook Salmon escapement estimates from weir counts on the Gisasa River through 2022 range from 503 to 4,023 fish per year. Summer Chum Salmon escapement estimates for the same period range from 3,300 to 261,305 fish per year. The Gisasa River weir study objectives for 2022 were to: (1) determine daily fish passage, estimate seasonal escapement, and describe run timing of adult Chinook and summer Chum Salmon, (2) determine age, sex, and length composition of adult Chinook and summer Chum Salmon, (3) document observations of resident fish, and (4) work with Tanana Chiefs Conference (TCC), as the tribal organization for the region, to transition operation of the project from USFWS staff to TCC.

Study Area

The Gisasa River headwaters originate in the Nulato Hills, and the river flows northeast as it passes through the Koyukuk National Wildlife Refuge. Approximately 112 km from its source, the Gisasa River enters the Koyukuk River at roughly 65° 15.149' N, 157° 42.925' W (USGS 1:63,360 series, Kateel River B-4 quadrangle), 90 km upriver from the confluence of the Koyukuk and Yukon rivers (Figure 1). Climate in the region is continental subarctic with dramatic seasonal temperature variations and low precipitation. Mean annual air temperature at the village of Galena, 64 km southeast of the Gisasa River is 3.8 °C with extremes ranging from 32 °C during summer to -57 °C during winter (USFWS 1993). The hydrology of this area is dynamic throughout the year, with lower flows generally occurring in late summer. Peak flows usually occur during spring break up or occasionally during summer high precipitation events. Rivers in the area generally begin to freeze during October and break up during May.

The weir site is located approximately 4 km upriver from the mouth of the Gisasa River. This section of the river is straight with generally laminar flow. The river channel cross section slopes gradually from both stream banks to the thalweg and is approximately 45 m wide. The water depth at the trap near the thalweg ranged from 66 to 96 cm (Appendix 2). Predominant substrate at the weir site consists of medium-sized gravel, 35 to 70 mm in diameter.

Methods

Weir Operation

A resistance board weir was used to enumerate and collect biological data from adult salmon as they migrated up the Gisasa River to spawn. The Gisasa River weir has been installed at the same site

since the project was initiated in 1994, following the construction and installation methods described by Tobin (1994). More detailed information on deployment of the Gisasa River weir can be found in Melegari and Wiswar (1995). A live trap was installed approximately mid-channel, near the thalweg, allowing fish to be recorded as they passed through the weir and, when necessary, the trap was closed to collect fish for sampling. The weir was visually inspected for integrity and cleaned of debris daily. Cleaning consisted of raking debris from the upstream surface of the weir or walking across each panel to submerge it enough to allow the current to wash debris downstream. The picket spacing within the trap and weir panels (3.5 cm between pickets) was narrow enough to prevent adult Chinook and summer Chum Salmon from passing through the weir. However, some individuals of smaller fish species, such as Arctic Grayling *Thymallus arcticus* and Whitefish *Coregonus* spp., were likely able to pass through the weir undetected.

A video camera system was installed on the upstream side of the weir trap box to capture video footage of migrating salmon and other species. The video camera box funneled fish into a narrow passage chute that allowed for the continual movement of fish through the weir, and motion capture video recording was used to count and identify all species passing through the weir. A new, narrower passage chute first deployed in 2015 helped improve species identification and made enumeration easier by limiting overlapping fish in the view window.

Once video counting began, motion capture features were enabled, and all counting was conducted from individual motion capture files. Motion capture files were saved to a hard drive and reviewed hourly. Fish were identified to species and total hourly counts were entered into an electronic data sheet daily. The video box was equipped with LED lights so that fish could be observed 24 hours per day. Adjustments to video settings and equipment were made as necessary to optimize image quality and performance of the system. During the first week of weir operation, motion capture performance was closely observed to verify that settings were adjusted properly and all fish passing the camera were detected. After the initial verification period, only motion capture video files were used for the remainder of the season. Accuracy was monitored throughout the season with short visual observation periods.

The target start date for weir operation is June 18. The end date of the project is determined inseason, when the daily count of both Chinook and summer Chum Salmon drops to less than 1% of the seasonal passage to date for three consecutive days, or when logistical constraints require stopping before this point is reached. A daily counting schedule was implemented, and fish were counted 24 hours per day. Escapement counts and sex ratios from the previous day were reported daily to the NAFWFO by satellite telephone.

Hydrological Data

Water depth (cm) was recorded twice daily at the weir trap at approximately 0800 hours and 2000 hours. Water temperature (°C) was taken twice daily with a handheld thermometer upstream of the weir in a section of river where water was well mixed.

Biological Data

Sex and length data and scale samples (to determine age) were collected from Chinook and summer Chum Salmon in a trap connected to the passage chute. Both ends of the trap were closed to retain fish during sampling but were open at all other times. To avoid temperature stress, sampling and fish handling were suspended if water temperatures remained above 17 °C for three consecutive days or any time temperatures reached or exceeded 20 °C (Shink 2020). The sample size goal for each

species was 220–240 fish for the season. This goal was based on a statistical calculation indicating that a minimum of 180 samples are required to obtain eight simultaneous 95% confidence intervals for age-sex classes such that each interval is no more than ± 0.1 unit wide (Bromaghin 1993). The sample size was increased by 25% above this minimum as an allowance for unreadable scales and other sampling errors. To accurately characterize the composition of the entire run, the total number of samples must be distributed throughout the run roughly in proportion to escapement, while not exceeding the target number of samples. As a pre-season guideline for sampling, the season was divided into three periods, or tertiles, based on historical fish passage counts, and the target sample size within each tertile was 73–80 fish. The tertile dates designated for Chinook Salmon were June 16–July 11, July 12–17, and July 18–31. The tertile dates designated for summer Chum Salmon were June 16–July 7, July 8–14, and July 15–31. To ensure consistent sampling effort towards meeting the target sample size for each tertile, daily and weekly targets were set. Salmon were sampled opportunistically during a day or week until these targets were met.

For age determination, three scales from each Chinook Salmon and one scale from each summer Chum Salmon were collected from the preferred sampling area (INPFC 1963). Scales were collected from the left side of the fish, two rows above the lateral line on a diagonal from the posterior insertion of the dorsal fin to the anterior insertion of the anal fin. Scales from both adult salmon species were sent to the Stock Biology group at ADF&G Division of Commercial Fisheries in Anchorage for age analysis (Eaton 2015). Fish length was measured from mid-eye to fork of tail (MEFL), and sex was visually determined from external characteristics such as kype development or the presence of an ovipositor. Previous studies have shown that nearly all age-1.1 and -1.2 Chinook Salmon, including those less than 650 mm (MEFL) are males (Brady 1983; Bales 2007; Karpovich and DuBois 2007). Therefore, all age-1.1 and -1.2 Chinook Salmon and those fish less than 650 mm (MEFL) were assumed to be males.

Data Analysis

The biological sampling guideline based on previous years' run timing is for crew guidance only and does not ensure that sample distribution over time will correspond to actual run timing. Therefore, age, sex, and length data were post-stratified during analysis by distributing them into two or more periods which had approximately equal fish passage numbers. Weighted estimates from each period were then combined to form seasonal estimates of age and sex composition.

Means and standard errors were calculated and reported for fish lengths within each age and sex class. Lengths within each age-sex class were assumed not to vary substantially over the seasonal progression of the run, so length calculations were made on pooled, unweighted samples within each age-sex class.

Age and sex composition was calculated as population proportions based on the number of sampled fish in each age and sex class. In each sampling period i , a number of fish n_i were sampled, and of these, a number n_{ij} were determined to be of class j (by age, sex, or age and sex combined). The proportion p_{ij} of fish of class j in sampling period i was estimated as:

$$\widehat{p}_{ij} = \frac{n_{ij}}{n_i} .$$

Sample proportions were weighted by the proportion N_i/N of the total population N available for sampling within period i (i.e., proportion of the total seasonal escapement counted during period i).

Weighted sample proportions were summed across K periods to provide a pooled seasonal estimate of the proportion of class j , as:

$$\hat{p}_j = \sum_{i=1}^K \frac{N_i}{N} \hat{p}_{ij}.$$

The variance of each seasonal age, sex, or age-sex class proportion was estimated as:

$$\widehat{\text{Var}}(\hat{p}_j) = \sum_{i=1}^K \left(1 - \frac{n_i}{N_i}\right) \left(\frac{N_i}{N}\right)^2 \frac{\hat{p}_{ij}(1 - \hat{p}_{ij})}{n_i - 1}.$$

Using the above equation for variance and assuming asymptotic normality, 95% confidence intervals (CI) were constructed as:

$$\hat{p}_j \pm 1.96 \sqrt{\widehat{\text{Var}}(\hat{p}_j)}.$$

Results and Discussion

Weir Operation

The target start date of June 18 was not met due to staffing shortages, and weir repair and maintenance issues. However, the weir was fully operational and fish tight at 2000 hours on June 29, with no fish passing through on that day. In 2022, the weir was installed in the same location as in 2019, angled slightly upstream on the left bank to ensure that the weir was oriented perpendicular to the flow of the main current. Counting continued throughout the season with no substantial interruptions. Weir operations were stopped for the season at 0900 hours on July 29. Counting ended prematurely this season due to concern of flash flooding. The video camera system worked well, and no fish were physically observed passing through the chute without detection. All fish passing through the weir were identified to species and enumerated.

Hydrological Data

The average river stage height during weir operations was 74 cm and ranged between 66 cm and 96 cm. Water temperature during the season averaged 15 °C and ranged from 11 °C to 20 °C (Figure 2, Appendix 2).

Biological Data

An estimated 503 Chinook Salmon and 3,300 summer Chum Salmon (Table 1) passed through the weir during the period of operation. The next most abundant species was Northern Pike *Esox lucius* ($N = 22$), followed by Longnose Sucker *Catostomus catostomus* ($N = 10$), Broad Whitefish *Coregonus nasus* ($N = 9$), Sockeye Salmon *O. nerka* ($N = 5$), Humpback Whitefish *Coregonus pidschian* ($N = 3$), Arctic Grayling *Thymallus arcticus* ($N = 1$), and Pink Salmon *O. gorbuscha* ($N = 1$).

Chinook Salmon — The estimated Chinook Salmon escapement ($N = 503$) was well below the 1995–2019 historical average annual escapement of 2,065 (Figure 3, Appendix 1). The first Chinook Salmon passed through the weir on June 30. Eight Chinook Salmon passed through the weir on the last day of counting, representing 1.5% of total escapement. We assume the estimated Chinook Salmon escapement to be an underestimate because a small number of Chinook Salmon likely passed the site after the weir was removed for the season. The mid-point of Chinook Salmon passage occurred on July 18, with the first and third quarter passage dates occurring on July 13 and July 22, respectively (Table 1, Figure 4). We assume the actual quartile passage dates would be later than reported here, as counting ended prematurely this season.

Sex and length data, and scale samples were collected from 42 Chinook Salmon. Due to the low total run size, there was difficulty obtaining samples, and the sample size goal was not achieved. Because of the insufficient number of samples, ASL analyses could not be performed for Chinook Salmon.

Summer Chum Salmon — The estimated summer Chum Salmon escapement ($N = 3,300$) was well below the 1995–2019 historical average annual escapement of 66,832 (Figure 5, Appendix 1). The first summer Chum Salmon passed through the weir on July 1. Four percent of total escapement ($n = 147$) passed through the weir on the last day of counting. We assume the estimated summer Chum Salmon escapement to be an underestimate because some summer Chum Salmon likely passed the site after the weir was removed for the season. The mid-point of summer Chum Salmon passage occurred on July 16, with the first and third quarter passage dates occurring on July 11 and July 24, respectively (Table 1, Figure 6). We assume the actual quartile passage dates would be later than reported here, as counting ended prematurely this season. The summer Chum Salmon escapement at the Gisasa River weir represented 2.1% of the entire Yukon River summer Chum Salmon run estimate generated by the Pilot Station sonar project (JTC 2022).

Sex and length data, and scale samples were collected from 294 summer Chum Salmon. Age was unable to be determined for 4.4% of the samples ($n = 13$), due to the scale being missing, inverted, illegible, regenerated, or absorbed (Table 2). Two age classes of summer Chum Salmon were identified, 0.3 and 0.4, from brood years 2018 and 2017, and each of these age classes comprised 50% of the escapement. Age classes 0.2 and 0.5 were not represented in Gisasa River samples in 2022 (Table 2).

The estimated sex composition for summer Chum Salmon was 55% female (Table 2). The overall female composition of 55% was almost equal to the 1995–2019 average of 52% female (Table 2, Appendices 3 and 4). Female summer Chum Salmon lengths ranged from 445 to 595 mm MEFL (Table 3). Male summer Chum Salmon lengths ranged from 450 to 615 mm MEFL (Table 3). Mean length-at-age of male summer Chum Salmon was larger than that of females for both age classes, 0.3 and 0.4 (Table 3).

The Gisasa River weir has produced 26 years of salmon escapement data (1994–2022, excluding 2018, 2020, and 2021). In addition to helping state and federal fisheries managers direct in-season management decisions, these data are instrumental to developing future run projections and evaluating escapement goals and harvest allocations throughout the Yukon River drainage. Long-term data sets such as these allow for the analysis of trends in salmon population status, size, length, age, and sex composition, which continues to be of great importance as stressors affecting Yukon River salmon, including climate change, disease, selective harvest, and overall demand, change over time.

Acknowledgments

Funding for this project was provided through the U.S. Fish and Wildlife Service, Office of Subsistence Management, Fisheries Resource Monitoring Program, under project 22-202. Appreciation is extended to the field crew, Eric Rowe, Erin McCarthy, and Scotty Starr; and to Sarah Conn for editorial review. We would also like to thank Courtney Berry and Jim O'Rourke of ADF&G, Commercial Fisheries Division, Arctic-Yukon-Kuskokwim Region, for their assistance with scale sample analysis, and to the City of Koyukuk, Koyukuk National Wildlife Refuge staff, and the Bureau of Land Management Alaska Fire Service in Galena, for logistical support.

References

- Bales, J. 2007. Salmon age and sex composition and mean lengths for the Yukon River area, 2005. Alaska Department of Fish and Game, Fishery Data Series No. 07-04, Anchorage.
- Barton, L. H. 1984. A catalog of Yukon River salmon spawning escapement surveys. Alaska Department of Fish and Game, Division of Commercial Fisheries. Fairbanks, Alaska.
- Brady, J. A. 1983. Lower Yukon River salmon test and commercial fisheries, 1981. Alaska Department of Fish and Game, Technical Data Report 89. 91 p.
- Bromaghin, J. F. 1993. Sample size determination for interval estimation of multinomial probabilities. *The American Statistician* 47:203–206.
- Eaton, S. E. 2015. Salmon age, sex, and length (ASL) sampling procedures for the Arctic-Yukon-Kuskokwim region. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 3A15-04, Anchorage.
- INPFC (International North Pacific Fisheries Commission). 1963. Annual report for 1961. Vancouver, British Columbia.
- JTC (Joint Technical Committee of the Yukon River U.S./Canada Panel). 2022. Yukon River salmon 2021 season summary and 2022 season outlook. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 3A22-01, Anchorage.
- Karpovich, S., and L. DuBois. 2007. Salmon age and sex composition and mean lengths for the Yukon River area, 2004. Alaska Department of Fish and Game, Fishery Data Series No. 07-05, Anchorage.
- Kruse, G. E. 1998. Salmon run failures in 1997–1998: A link to anomalous ocean conditions? *Alaska Fisheries Resource Bulletin* 5(1):55–63.
- Melegari, J. L., and D. W. Wiswar. 1995. Abundance and run timing of adult salmon in the Gisasa River, Koyukuk National Wildlife Refuge, Alaska, 1994. U.S. Fish and Wildlife Service, Fairbanks Fishery Resource Office, Fishery Data Series Number 95-1, Fairbanks, Alaska.
- Shink, K. G. 2020. Guidelines for handling and sampling adult salmon during periods of elevated water temperatures in Alaskan rivers. Alaska Fisheries Data Series No. 2020-1, U.S. Fish and Wildlife Service, Anchorage.
- Tobin, J. H. 1994. Construction and performance of a portable resistance board weir for counting migrating adult salmon in rivers. U.S. Fish and Wildlife Service, Kenai Fishery Resource Office, Alaska Fisheries Technical Report Number 22, Kenai, Alaska.
- USFWS (U.S. Fish and Wildlife Service). 1993. Fishery management plan–Koyukuk National Wildlife Refuge. Fairbanks Fishery Resource Office, Fairbanks, Alaska.
- Vania, T., V. Golembeski, B. M. Borba, T. L. Lingnau, J. S. Hayes, K. R. Boeck, and W. H. Busher. 2002. Annual management report Yukon and Northern areas 2000. Alaska Department of Fish and Game, Regional Information Report Number 3A02-29, Anchorage, Alaska.

Table 1. — Daily and cumulative (cum) estimates of Chinook Salmon and summer Chum Salmon passage, and daily estimates of other species, at the Gisasa River weir, Alaska, 2022.

Date	Chinook Salmon daily	Chinook Salmon cum	Chum Salmon daily	Chum Salmon cum	Sockeye Salmon daily	Northern Pike daily	Arctic Grayling daily	Whitefish daily	Longnose Sucker daily	Pink Salmon daily
Jun-29	0	0	0	0	0	0	0	0	0	0
Jun-30	1	1	0	0	0	1	0	0	0	0
Jul-01	0	1	1	1	0	0	0	1	0	0
Jul-02	0	1	8	9	0	0	1	2	0	0
Jul-03	0	1	20	29	0	0	0	0	0	0
Jul-04	2	3	28	57	0	1	0	1	0	0
Jul-05	2	5	20	77	0	3	0	1	0	0
Jul-06	3	8	34	111	0	1	0	0	0	0
Jul-07	12	20	39	150	0	1	0	0	0	0
Jul-08	12	32	76	226	0	1	0	1	1	0
Jul-09	8	40	201	427	0	0	0	0	2	0
Jul-10	15	55	292	719	0	1	0	0	2	0
Jul-11	42	97	291	1,010*	0	1	0	0	1	0
Jul-12	23	120	162	1,172	0	2	0	0	0	0
Jul-13	19	139*	128	1,300	0	0	0	0	1	0
Jul-14	29	168	233	1,533	0	1	0	0	0	0
Jul-15	16	184	108	1,641	0	1	0	0	0	0
Jul-16	23	207	98	1,739*	0	3	0	0	0	0
Jul-17	30	237	124	1,863	0	0	0	0	0	0
Jul-18	51	288*	120	1,983	0	0	0	1	0	0
Jul-19	6	294	50	2,033	0	0	0	0	0	0
Jul-20	10	304	58	2,091	1	0	0	0	0	0
Jul-21	71	375	101	2,191	1	1	0	0	0	0
Jul-22	35	410*	77	2,269	0	0	0	0	0	0
Jul-23	11	421	49	2,318	0	0	0	1	0	0
Jul-24	7	428	206	2,524*	1	1	0	0	1	0
Jul-25	21	449	261	2,785	0	1	0	3	0	0
Jul-26	30	479	152	2,937	1	1	0	0	1	0
Jul-27	8	487	92	3,029	1	0	0	0	0	0
Jul-28	8	495	124	3,153	0	1	0	0	0	0
Jul-29	8	503	147	3,300	0	0	0	1	1	1
Total	503	503	3,300	3,300	5	22	1	12	10	1

Note: Asterisks (*) indicate first, mid, and third quarter points of Chinook Salmon and summer Chum Salmon passage estimates.

Table 2. — Age and sex ratio estimates of summer Chum Salmon at the Gisasa River weir, Alaska, 2022.

Strata dates	Run size (N)	Sample size (n)	% Female	Unknown age	Brood year 2019 (age 0.2)	Brood year 2018 (age 0.3)	Brood year 2017 (age 0.4)	Brood year 2016 (age 0.5)
July 04–13	1,300	53	62%	9	0% (0.0)	45% (7.6)	55% (7.6)	0% (0.0)
July 14–17	563	97	56%	11	0% (0.0)	53% (5.4)	47% (5.4)	0% (0.0)
July 18–24	661	70	51%	4	0% (0.0)	47% (6.2)	53% (6.2)	0% (0.0)
July 25–28	776	74	45%	12	0% (0.0)	56% (6.4)	44% (6.4)	0% (0.0)
Total	3,300	294	55%	36	0% (0.0)	50% (3.7)	50% (3.7)	0% (0.0)
Female	1,809	156	-	8	0% (0.0)	50% (5.0)	50% (5.0)	0% (0.0)
Male	1,491	125	-	28	0% (0.0)	48% (5.6)	52% (5.6)	0% (0.0)

Note: Standard errors are in parenthesis. Season totals calculated from weighted strata totals. Unknown age indicates numbers of fish that could not be aged from the scales sampled and were not included in age calculations.

Table 3. — Length-at-age of male and female summer Chum Salmon sampled at Gisasa River weir, Alaska, 2022.

Age	Male N	Male mean	Male SE	Male median	Male range	Female N	Female mean	Female SE	Female median	Female range
0.2	-	-	-	-	-	-	-	-	-	-
0.3	52	525	29.3	528	450–585	79	511	25.9	510	445–595
0.4	56	546	28.8	545	490–615	69	527	27.5	525	460–595
0.5	-	-	-	-	-	-	-	-	-	-
Total	108	536	30.8	535	450–615	148	518	27.6	515	445–595

Note: Length was measured mid-eye to fork, MEFL (mm).




Figure 1. — Location of the Gisasa River weir and other active and historical tributary escapement project sites in the Koyukuk River drainage, Alaska.




Figure 2. — Average daily river stage height and water temperature at the Gisasa River weir, Alaska, 2022.

Note: Horizontal lines represent the 2022 average river stage height (dashed) and water temperature (solid). The average was calculated from the morning and evening water temperature and river stage height values.




Figure 3. — Chinook Salmon escapement estimates at the Gisasa River weir, Alaska, 1994–2022.

Note: *Data from the first year of operation (1994) represent only a partial count; counting did not begin until July 10, after the run was underway. These data are not included in the average. **Data from 2014 represent a partial count due to high water events and are not included in the average. The horizontal line represents the 1995–2019 average. The weir was not operated in 2018, 2020, and 2021.




Figure 4. — Daily 2022 and average daily (1995–2021) Chinook Salmon passage estimates through the Gisasa River weir, Alaska.

Note: The weir did not operate in 2018, 2020, and 2021*.




Figure 5. — Summer Chum Salmon escapement estimates at the Gisasa River weir, Alaska, 1994–2022.

Note: *Data from the first year of operation (1994) represent only a partial count; counting did not begin until July 10, after the run was underway. These data are not included in averages. **Data from 2014 represent a partial count due to high water events and are not included in the average. The horizontal lines represent the 1995–2019 average (solid) and median (dashed). The weir was not operated in 2018, 2020, and 2021.




Figure 6. — Daily 2022 and average daily (1995–2019) Summer Chum Salmon passage estimates through the Gisasa River weir, Alaska.

Note: The weir did not operate in 2018, 2020, and 2021*.

Appendix 1. — Historical Chinook Salmon and Summer Chum Salmon escapement estimates in the Gisasa River, Alaska, 1960–2022.

Year	Aerial index estimates - Chinook Salmon	Aerial index estimates - summer Chum Salmon	Aerial survey rating	Weir estimates - Chinook Salmon	Weir estimates - summer Chum Salmon
1960	300	400	Good		
1961	266	0	Good		
1974	161	22,022	Good		
1975	385	56,904	Good		
1976	332	21,342	Good		
1977	255	2,204	Good		
1978	45	9,280	Good		
1979	484	10,962	Good		
1980	951	10,388	Good		
1982	421	33	Good		
1983	572	2,356	Good		
1985	735	13,232	Good		
1986	1,346	12,114	Good		
1987	731	2,123	Good		
1988	797	9,284	Good		
1990	884	450	Good		
1991	1,690	7,003	Good		
1992	910	9,300	Good		
1993	1,573	1,581	Good		
1994	2,775	6,827	Good	2,888	51,116
1995	410	6,458	Good	4,023	136,886
1996				1,991	158,752
1997	144	686	Good	3,764	31,800
1998	889		Poor	2,414	21,142
1999				2,644	10,155
2000				2,089	11,410
2001	1,298		Good	3,052	17,946
2002	506		Good	2,025	33,481
2003				1,901	25,999
2004	731		Good	1,774	37,851
2005	958		Good	3,111	172,259
2006	843	1,000	Fair	3,031	261,306
2007	593		Fair	1,427	46,257
2008	487	20,470	Fair	1,738	36,938
2009	515	1,060	Good	1,955	25,904
2010	564	1,096	Fair	1,516	47,669
2011	906	13,228	Good	2,692	95,796
2012				1,323	83,423
2013	201	9,300	Surveyed too late	1,126	80,055
2014				1,589	32,523
2015	558	5,601	Good	1,319	42,747
2016				1,395	66,670
2017	452	8,058	Fair	1,083	73,584

-continued-

Appendix 2. — Page 2 of 2.

Year	Aerial index estimates - Chinook Salmon	Aerial index estimates - summer Chum Salmon	Aerial survey rating	Weir estimates - Chinook Salmon	Weir estimates - summer Chum Salmon
2018				ND ^a	ND ^a
2019				1,328	19,099
2020				ND ^b	ND ^b
2021				ND ^b	ND ^b
2022				503	3,300

Note: Aerial index data are from Barton (1984) and the Alaska Department of Fish and Game Joint Technical Committee (JTC) (2022).

^a No escapement estimates due to persistent flooding.



^b No escapement estimates due to COVID-19 pandemic.

Appendix 2. — Water depth, water temperature, and air temperature data collected at the Gisasa River weir, Alaska, 2022.

Date	AM water depth (cm)	PM water depth (cm)	AM water temp (°C)	PM water temp (°C)	Low air temp (°C)	High air temp (°C)
27-Jun	74	74	14	15	-	-
28-Jun	74	74	14	15	-	-
29-Jun	74	74	14	15	-	-
30-Jun	74	74	14	17	-	-
1-Jul	72	71	14	18	-	-
2-Jul	70	70	15	18	-	-
3-Jul	72	72	15	18	13	20
4-Jul	74	75	16	16	14	20
5-Jul	76	76	15	16	11	20
6-Jul	74	74	14	18	9	24
7-Jul	74	74	16	17	13	22
8-Jul	72	71	16	19	9	27
9-Jul	72	71	16	19	11	28
10-Jul	70	70	17	20	12	27
11-Jul	68	68	16	19	12	22
12-Jul	67	68	16	16	10	17
13-Jul	68	68	14	17	6	21
14-Jul	68	67	15	18	5	22
15-Jul	67	66	15	14	7	12
16-Jul	67	67	13	15	10	19
17-Jul	68	68	14	16	6	21
18-Jul	68	68	15	15	4	14
19-Jul	67	66	12	13	5	14
20-Jul	66	68	12	12	6	12
21-Jul	70	80	11	14	7	14
22-Jul	95	91	11	13	3	18
23-Jul	85	83	11	12	6	18
24-Jul	80	78	11	14	9	15
25-Jul	78	80	11	14	4	14
26-Jul	85	94	11	12	-	-
27-Jul	96	-	12	-	-	-
Average	74	73	14	16	8	19

Note: Water depth is the water level at the trap.

Appendix 3. — Historical percentages of female Chinook Salmon and summer Chum Salmon for the Gisasa River weir, Alaska, 1994–2022.



Note: *Data from the first year of operation (1994) represent only a partial count; counting did not begin until July 10, after the run was underway and these data are not included in the average. **Data from 2014 represent a partial count due to high water events and are not included in the average. The horizontal line represents the 1995–2019 average. The weir was not operated in 2018, 2020, and 2021. The ASL sample size goal was not met for Chinook Salmon in 2022, and therefore data were not analyzed.

Appendix 4. — Historical percentages of female Chinook Salmon and summer Chum Salmon sampled at the Gisasa River weir, Alaska.

Year	Chinook % female	Summer Chum % female
*1994	39	64
1995	44	46
1996	20	51
1997	23	51
1998	17	46
1999	29	52
2000	30	50
2001	42	49
2002	21	48
2003	35	48
2004	31	44
2005	33	45
2006	32	55
2007	34	57
2008	18	49
2009	26	53
2010	24	53
2011	17	54
2012	37	56
2013	31	56
**2014	18	51
2015	28	55
2016	28	58
2017	28	57
2018		
2019	30	59
2020		
2021		
2022		53
Average	28	52

Note: *Indicates incomplete data from the first year of operation (1994); data collection did not begin until July 10.
 **Indicates incomplete data from 2014 when high water events impeded weir operation for most of the season. The weir was not operated in 2018, 2020, and 2021. The ASL sample size goal was not met for Chinook Salmon in 2022, and therefore data were not analyzed.