

# *Pristobrycon striolatus* (a piranha, no common name)

## Ecological Risk Screening Summary

U.S. Fish & Wildlife, May 2012  
Revised, November 2018  
Web Version, 1/25/2021

Organism Type: Fish  
Overall Risk Assessment Category: Uncertain




Image: Albert C. L. G. Gunther. Public Domain, hosted by Freshwater and Marine Image Bank. Available: <https://eol.org/media/9336782>. (November 7, 2018).

## 1 Native Range and Status in the United States

---

### Native Range

From Froese and Pauly (2018):

“South America: Amazon and Orinoco River basins [Brazil, Venezuela] and north and eastern Guiana Shield rivers [French Guiana, Guyana, Suriname].”

“[In Brazil:] Known from the Amazon River basin [Jégu 2003]. Occurs in Para and Serra do Roncador [Planquette et al. 1996].”

“[In French Guiana:] Present in the Maroni River [Planquette et al. 1996].”

From Diaz-Sarmiento and Alvares-León (2003):

“*Serrasalmus (Pygocentrus) nattereri*, *Serrasalmus rhombeus*, *Serrasalmus spilopleura*, and *Serrasalmus [Pristobrycon] striolatus* have been reported from the Caquetá River [Colombia].”

## **Status in the United States**

No records of *Pristobrycon striolatus* in the wild or in trade in the United States were found.

The Florida Fish and Wildlife Conservation Commission has listed the fish, *Pristobrycon striolatus* as a prohibited species. Prohibited nonnative species (FFWCC 2018), "are considered to be dangerous to the ecology and/or the health and welfare of the people of Florida. These species are not allowed to be personally possessed or used for commercial activities."

*Pristobrycon striolatus* is a prohibited species in Alabama (Alabama DCNR 2019). “No person, firm, corporation, partnership, or association shall possess, sell, offer for sale, import, bring, release, or cause to be brought or imported into the State of Alabama” a prohibited species.

Piranhas are prohibited species in Arkansas (Arkansas GFC 2019). *Pristobrycon striolatus* is a species of piranha.

Species of piranha (including *Pristobrycon striolatus*) are prohibited as pets in Georgia (Georgia DNR 2020).

Possession of species of piranha (including *Pristobrycon striolatus*) is prohibited in Louisiana (Louisiana State Legislature 2019).

*Pristobrycon striolatus* falls within Group IV of New Mexico’s Department of Game and Fish Director’s Species Importation List (New Mexico Department of Game and Fish 2010). Group IV species “are prohibited for the general public but may be allowed for, scientific study, department approved restoration and recovery plans, zoological display, temporary events/entertainment, use as service animal or by a qualified expert.”

*Pristobrycon striolatus* is a prohibited species in Texas (Texas Parks and Wildlife 2020).

From Virginia DWR (2020):

“A special permit is required, and may be issued by the Department, if consistent with the Department’s fish and wildlife management program, to import, possess, or sell the following non-native (exotic) amphibians, fish, mollusks, aquatic invertebrates, and reptiles: [...] piranhas,”

## **Means of Introductions in the United States**

No records of *Pristobrycon striolatus* in the wild in the United States were found.

## Remarks

Information searches were conducted using the valid name *Pristobrycon striolatus* and the synonym *Serrasalmus striolatus*.

## 2 Biology and Ecology

---

### Taxonomic Hierarchy and Taxonomic Standing

According to Fricke et al. (2018), *Pristobrycon striolatus* is the valid name for this species. It was originally described as *Serrasalmo (Pygocentrus) striolatus* and has been previously known as *Serrasalmus striolatus*.

From ITIS (2018):

Kingdom Animalia

Subkingdom Bilateria

Infrakingdom Deuterostomia

Phylum Chordata

Subphylum Vertebrata

Infraphylum Gnathostomata

Superclass Actinopterygii

Class Teleostei

Superorder Ostariophysi

Order Characiformes

Family Characidae

Genus *Pristobrycon*

Species *Pristobrycon striolatus* (Steindachner, 1908)

### Size, Weight, and Age Range

From Froese and Pauly (2018):

“Max length : 20.1 cm TL male/unsexed; [Giarrizzo et al. 2015]; max. published weight: 230.00 g [Giarrizzo et al. 2015]”

### Environment

From Froese and Pauly (2018):

“Freshwater; benthopelagic.”

### Climate

From Froese and Pauly (2018):

“Tropical”

## **Distribution Outside the United States**

### **Native**

From Froese and Pauly (2018):

“South America: Amazon and Orinoco River basins [Brazil, Venezuela] and north and eastern Guiana Shield rivers [French Guiana, Guyana, Suriname].”

“[In Brazil:] Known from the Amazon River basin [Jégu 2003]. Occurs in Para and Serra do Roncador [Planquette et al. 1996].”

“[In French Guiana:] Present in the Maroni River [Planquette et al. 1996].”

From Diaz-Sarmiento and Alvares-León (2003):

“*Serrasalmus (Pygocentrus) nattereri*, *Serrasalmus rhombeus*, *Serrasalmus spilopleura*, and *Serrasalmus [Pristobrycon] striolatus* have been reported from the Caquetá River [Colombia].”

### **Introduced**

No records of *Pristobrycon striolatus* introductions were found.

## **Means of Introduction Outside the United States**

No records of *Pristobrycon striolatus* introductions were found.

## **Short Description**

A short description of *Pristobrycon striolatus* was not found.

## **Biology**

From Froese and Pauly (2018):

“Prefers to inhabit the upper reaches of main rivers. Stomach content analysis showed that its food consisted mainly of seeds and to a lesser degree, morsels of fins [Planquette et al. 1996].”

## **Human Uses**

From Froese and Pauly (2018):

“Rarely captured.”

According to Begossi et al. (2019), *Pristobrycon striolatus* is consumed as a food fish in coastal and inland areas of Brazil.

## Diseases

No records of OIE-reportable diseases (OIE 2021) were found for *Pristobrycon striolatus*.

Casal et al. (1997) described *Henneguya striolata* as a parasite of *P. striolatus* (under the name *Serrasalmus striolatus*).

Brandão et al. (2013) list *P. striolatus* as a host for *Anacanthorus amazonicus*, *Anacanthorus cinctus*, *Anacanthorus crytocaulus*, and *Anacanthorus lasiophallus*.

Kritsky et al. (1997) list *P. striolatus* as a host for *Amphithecium prodotum*, *Pithanothecium amazonensis*, and *Pithanothecium piranhus*.

Kritsky et al. (1996) list *P. striolatus* as a host for *Notozothecium robustum*.

Kritsky et al. (1998) list *P. striolatus* as a host for *Enallothecium variabilum*.

Whipps et al. (2012) list *P. striolatus* (as *Serrasalmus striolatus*) as a host for *Calyptospora spinosa*.

Poelen et al. (2014) list *Anavilhanatrema* sp. as an additional parasite of *P. striolatus*.

## Threat to Humans

From Froese and Pauly (2018):

“Harmless”

## 3 Impacts of Introductions

---

No records of *Pristobrycon striolatus* introductions were found; therefore, there is no information on impacts of introduction to evaluate.

*P. striolatus* is regulated in multiple States.

## 4 History of Invasiveness

---

No records of introductions of *Pristobrycon striolatus* were found, therefore the history of invasiveness is classified as “no known nonnative population.”

## 5 Global Distribution

---



**Figure 1.** Known global distribution of *Pristobrycon striolatus*. Locations are in Colombia, Venezuela, French Guiana, Guyana, Suriname, and Brazil. Map from GBIF Secretariat (2018). The location in southern Brazil was not used to select source points for the climate match. The recorded collection point does not match the recorded coordinates (GBIF Secretariat 2018).

## 6 Distribution Within the United States


---

No records of *Pristobrycon striolatus* in the wild in the United States were found.


# 7 Climate Matching

## Summary of Climate Matching Analysis

The climate match for *Pristobrycon striolatus* was mostly low for the contiguous United States. There were some areas of medium match along the Gulf Coasts of Texas and Louisiana. Southern Florida had a medium match with the very tip of the state having a high match. The Climate 6 score (Sanders et al. 2018; 16 climate variables; Euclidean distance) for contiguous United States was 0.001, low (scores between 0.000 and 0.005, inclusive, are classified as low). All States had low individual climate 6 scores except for Florida which had a medium individual score.



**Figure 2.** RAMP (Sanders et al. 2018) source map showing weather stations in South America selected as source locations (red; Colombia, Venezuela, French Guiana, Guyana, Suriname, Brazil, Peru) and non-source locations (gray) for *Pristobrycon striolatus* climate matching. Source locations from GBIF Secretariat (2018). Selected source locations are within 100 km of one or more species occurrences, and do not necessarily represent the locations of occurrences themselves.



**Figure 3.** Map of RAMP (Sanders et al. 2018) climate matches for *Pristobrycon striolatus* in the contiguous United States based on source locations reported by GBIF Secretariat (2018). Counts of climate match scores are tabulated on the left. 0/Blue = Lowest match, 10/Red = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: (Count of target points with climate scores 6-10)/ (Count of all target points)	Overall Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
$\geq 0.103$	High

## 8 Certainty of Assessment

The certainty of assessment for *Pristobrycon striolatus* is low. There is minimal information available for this species. No records of introduction were found, therefore there is no information on impacts of introduction available to evaluate.



## 9 Risk Assessment

---

### Summary of Risk to the Contiguous United States

*Pristobrycon striolatus* is a species of piranha native to Atlantic river basins in South America from the Orinoco to the Amazon. It eats seeds and fruits as well as fish fins and is used as a food source in Brazil. The history of invasiveness is classified as “no known nonnative population.” No records of introduction were found. *P. striolatus* is regulated in multiple States. The climate match was low. There were small areas of medium match around the Gulf Coast along with medium matches for southern Florida and a high match at the very southern tip of Florida. The certainty of assessment is low. The overall risk assessment is uncertain.

### Assessment Elements

- **History of Invasiveness (Sec. 4): No Known Nonnative Population**
- **Overall Climate Match Category (Sec. 7): Low**
- **Certainty of Assessment (Sec. 8): Low**
- **Remarks/Important additional information: No additional information**
- **Overall Risk Assessment Category: Uncertain**

## 10 Literature Cited

---

**Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 11.**

- Alabama [DCNR] Department of Conservation and Natural Resources. 2019. Restrictions on possession, sale, importation and/or release of certain animals and fish. Alabama Department of Conservation and Natural Resources Administrative Code, Chapter 220-2-.26.
- Arkansas [GFC] Game and Fish Commission. 2019. Certain exotic species prohibited. Arkansas Game and Fish Commission Code of Regulations 26.13.
- Begossi A, Salivonchyk SV, Hallwass G, Hanazaki N, Lopes PFM, Silvano RAM, Dumaresq D, Pittock J. 2019. Fish consumption on the Amazon: a review of biodiversity, hydropower and food security issues. *Brazilian Journal of Biology* 79(2):345–357.
- Brandão H, Yamada FH, de Melo Toledo G, Carvalho ED, da Silva RJ. 2013. Monogeneans (Dactylogyridae) parasitizing gills of *Salminus hilarii* from a Neotropical reservoir, Brazil. *Brazilian Journal of Veterinary Parasitology* 22:579–587.
- Casal G, Matos E, Azevedo C. 1997. Some ultrastructural aspects of *Henneguya striolata* sp. nov. (Myxozoa, Myxosporea), a parasite of the Amazonian fish *Serrasalmus striolatus*. *Parasitology Research* 83:93–95.

- Diaz-Sarmiento JA, Alvarez-León R. 2003. Migratory fishes of the Colombian Amazon. Pages 303–344 in Carolsfeld J, Harvey B, Ross C, Baer A, editors. Migratory fishes of South America. Ottawa, Canada: The World Bank, International Development Research Centre.
- [FFWCC] Florida Fish and Wildlife Conservation Commission. 2018. Prohibited species list. Tallahassee, Florida: Florida Fish and Wildlife Conservation Commission. Available: <http://myfwc.com/wildlifehabitats/nonnatives/regulations/prohibited/> (November 2018).
- Fricke R, Eschmeyer WN, van der Laan R, editors. 2018. Catalog of fishes: genera, species, references. California Academy of Science. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (November 2018).
- Froese R, Pauly D, editors. 2018. *Pristobrycon striolatus* (Steindachner, 1908). FishBase. Available: <http://www.fishbase.org/summary/Pristobrycon-striolatus.html> (November 2018).
- GBIF Secretariat. 2018. GBIF backbone taxonomy: *Pristobrycon striolatus* (Steindachner, 1908). Copenhagen: Global Biodiversity Information Facility. Available: <https://www.gbif.org/species/2354222> (November 2018).
- Georgia [DNR] Department of Natural Resources. 2020. Wild animals/exotics. Social Circle, Georgia: Georgia Department of Natural Resources Law Enforcement Division. Available: <http://gadnrle.org/exotics> (November 2020).
- [ITIS] Integrated Taxonomic Information System. 2018. *Pristobrycon striolatus* (Steindachner, 1908). Reston, Virginia: Integrated Taxonomic Information System. Available: [https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=641596#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=641596#null) (November 2018).
- Kritsky DC, Boeger WA, Jégu M. 1996. Neotropical Monogenoidea. 28. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil and French Guiana: Species of *Notozothecium* Boeger and Kritsky, 1988, and *Mymarothecium* gen. n. Journal of Helminthological Society of Washington 63:153–175.
- Kritsky DC, Boeger WA, Jégu M. 1997. Neotropical Monogenoidea. 29. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil: species of *Amphithecium* Boeger and Kritsky, 1988, *Heterothecium* gen. n. and *Pithanothecium* gen. n. Journal of Helminthological Society of Washington 64:25–54.
- Kritsky DC, Boeger WA, Jégu M. 1998. Neotropical Monogenoidea. 31. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil: Species of *Notothecium* Boeger and Kritsky, 1988, and *Enallothecium* gen. n. Journal of Helminthological Society of Washington 65:31–49.

- Louisiana State Legislature. 2019. Exotic fish; importation, sale, and possession of certain exotic species prohibited; permit required; penalty. Louisiana Revised Statutes, Title 56, Section 319.
- New Mexico Department of Game and Fish. 2010. Director’s species importation list. Santa Fe, New Mexico: New Mexico Department of Game and Fish. Available: [http://www.wildlife.state.nm.us/download/enforcement/importation/information/Directors-Species-Importation-List-08\\_03\\_2010.pdf](http://www.wildlife.state.nm.us/download/enforcement/importation/information/Directors-Species-Importation-List-08_03_2010.pdf) (November 2020).
- [OIE] World Organisation for Animal Health. 2021. OIE-listed diseases, infections and infestations in force in 2020. Available: <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2021/> (January 2021).
- Poelen JH, Simons JD, Mungall CJ. 2014. Global Biotic Interactions: an open infrastructure to share and analyze species-interaction datasets. *Ecological Informatics* 24:148–159.
- Sanders S, Castiglione C, Hoff M. 2018. Risk Assessment Mapping Program: RAMP. Version 3.1. U.S. Fish and Wildlife Service.
- Texas Parks and Wildlife. 2020. Invasive, prohibited and exotic species. Austin, Texas: Texas Parks and Wildlife. Available: [https://tpwd.texas.gov/huntwild/wild/species/exotic/prohibited\\_aquatic.phtml](https://tpwd.texas.gov/huntwild/wild/species/exotic/prohibited_aquatic.phtml) (November 2020).
- Virginia [DWR] Department of Wildlife Resources. 2020. Nongame fish, reptile, amphibian and aquatic invertebrate regulations. Henrico, Virginia: Virginia Department of Wildlife Resources. Available: <https://dwr.virginia.gov/fishing/regulations/nongame/> (November 2020).
- Whipps CM, Fournie JW, Morrison DA, Azevedo C, Matos E, Thebo P, Kent ML. 2012. Phylogeny of fish-infecting *Calyptospora* species (Apicomplexa: Eimeriorina). *Parasitological Research* 111:1331–1342.

## 11 Literature Cited in Quoted Material

---

**Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.**

- Giarrizzo T, de Sena Oliveira RR, Andrade MC, Gonçalves AP, Barbosa TAP, Martins AR, Marques DK, dos Santos JLB, da S. Frois R de P, de Albuquerque TPO, de A. Montag LF, Camargo M, de Sousa LM. 2015. Length-weight and length-length relationships for 135 fish species from the Xingu River (Amazon basin, Brazil). *Journal of Applied Ichthyology* 31:514–424.

Jégu M. 2003. Serrasalminae (Pacus and piranhas). Pages 182–196 *in* Reis RE, Kullander SO, Ferraris CJ Jr, editors. Checklist of the freshwater fishes of South and Central America. Porto Alegre, Brazil: EDIPUCRS.

Planquette P, Keith P, Le Bail P-Y. 1996. Atlas des poissons d'eau douce de Guyane. Tome 1. Collection du Patrimoine Naturel volume 22. Paris: MNHN and INRA.