



## Klamath River Outmigrant Monitoring Update — June 6, 2022

Synopsis: The outmigration of juvenile salmonids is monitored annually on the mainstem Klamath River by the USFWS Arcata Fish and Wildlife Office (AFWO), the Karuk Tribe of California, and the Yurok Tribe of California. The objectives of this collaborative project are to:

1. Estimate the weekly abundance of juvenile Chinook Salmon and collect pertinent biological data such as fork length and presence of clinical signs of disease at four selected locations on the mainstem Klamath River.
2. Examine subsamples of Chinook Salmon, Coho Salmon, and steelhead for external disease indicators and collect, preserve, and deliver weekly-stratified, random samples of young-of-the-year (YOY) Chinook Salmon to the Service's California–Nevada Fish Health Center (CA–NV FHC) for conducting qPCR assays to estimate *Ceratonova shasta* infection rate in the outmigrant population.
3. Collect relative abundance and biological data on Coho Salmon and steelhead at the four locations on the mainstem Klamath River.

Information generated by this study are used for a variety of purposes, including stock-recruitment analyses, to inform flow management decisions, to further refine a fish disease model, and to validate and calibrate the S3 (Stream Salmonid Simulator) Chinook Salmon production model, among others.

Monitoring is conducted at four sites on the mainstem Klamath River between Iron Gate Dam (IGD; rkm 309.65) and the Trinity River confluence (rkm 64.3). The upstream-most site (rkm 307.75), referred to as the 'Bogus Trap Site' is located on the right bank downstream of the Bogus Creek confluence on Blue Heron RV Park property. The second location is the 'I-5 Trap Site' (rkm 293.55), which is positioned on the left bank downstream of the Carson Creek confluence and upstream of the I-5 bridge river crossing. The 'Kinsman Trap Site' (rkm 237.55) is positioned in a side channel on the left bank just upstream of the Kinsman Creek confluence. The 'Weitchpec Trap Site' (rkm 65) is the farthest downstream and is 0.7 km upstream of the Trinity River confluence behind the Yurok Tribal office in Weitchpec, California.

Trapping at the Bogus Trap Site is conducted using a single 3.1-m wide and 1.6-m tall frame net. Sampling at the I-5 Trap Site is conducted using two in-line 8-ft diameter rotary screw traps (RST) and one 3.1-m by 1.6-m frame net. One 5-ft diameter RST is used to capture fish at the Kinsman Trap Site. The Weitchpec trap site uses one 8-ft diameter RST on the south bank and one to two 3.1-m by 1.6-m frame nets on the north bank. Traps are typically operated four nights per week (Monday through Thursday) and checked once daily while in operation. Trapping began the week of February 28 [Calendar Week (CW) 10] at all sites in 2022.

This project update provides an in-season summary of the total catch (Table 1) and mean catch- per-day by week (Table 2) of Chinook Salmon, Coho Salmon, and steelhead at each trap site. In addition, we provide weekly estimates of mean fork length of YOY Chinook and Coho salmon from each of the four trap sites (Table 3). Expansions to generate weekly-stratified abundance estimates are calculated after the end of the sampling season and are not presented

here. Trap efficiency, a measure of the proportion of fish moving past a trap site that are caught, varies weekly. *Therefore, raw catch numbers are not representative of actual abundance and we advise against using weekly raw catch numbers to make inferences on temporal abundance.*

Included in this project update is a weekly-stratified summary of clinical signs of disease observed in the catch for the trap and seine sites (Table 4). Note that these data are based on the visual presence of external symptoms of disease, which may not always be revealed by infected fish. The percentage of live YOY Chinook Salmon in the trap and seine catches that exhibit distended bellies, gill fungus, and pale gills are presented separately for each site on a weekly basis (Table 4). Distended bellies may be a clinical sign of infection by the myxosporean parasites, *Ceratonova shasta* and *Parvicapsula minibicornis*. Gills of juvenile salmonids  $\geq 45$  mm FL are evaluated for color (red, pale/pink, white, or tan) and condition (normal, eroded, or fungal). Pale gills may be due to anemia associated with *P. minibicornis* infection. Gill fungus is likely *Saprolegnia* growing upon a columnaris (*Flavobacterium columnare*) infection.

To determine infection rates more accurately for the outmigrant juvenile Chinook Salmon population passing the Kinsman Trap Site, I-5 Trap Site, and Weitchpec Trap Site, weekly-stratified random samples are collected, preserved, and delivered to the CA–NV FHC to process using qPCR assays. This season's fish health sampling will begin the week of March 20. The CA–NV FHC investigates infection rates of *C. shasta* and other pathogens in juvenile salmonids in the Klamath River annually. The CA–NV FHC releases regular updates (which are available on the [USFWS online library](#)) and a final report for each season.

We also present daily mean discharge below IGD (Figure 1), at the Kinsman Trap Site (Figure 2), and at the Weitchpec Trap Site (Figure 3) from late February to July to help portray pertinent flow conditions. Discharge at the Bogus and I-5 trap sites are represented by USGS Gauging Station 11516530 (Klamath River below IGD, California). Discharge at USGS 11520500 (Klamath River near Seiad Valley, California) minus discharge at USGS 11519500 (Scott River near Fort Jones, California) is used as a surrogate flow for the Kinsman Trap Site. Discharge at USGS 10523000 (Klamath River at Orleans, California) is used to represent flow at the Weitchpec Trap Site.

If you have any questions regarding this summary, please contact Tyler Wallin (tyler\_wallin@fws.gov) or Bill Pinnix (bill\_pinnix@fws.gov).

Table 1. In-season summary of the total catch by week of adipose fin-clipped (AD Clip) and non-adipose fin-clipped (No Clip) Chinook Salmon and steelhead and left maxillary-clipped (LM Clip) and non-maxillary clipped (No Clip) Coho Salmon by trap at the Bogus, I-5, and Kinsman trap sites on the mainstem Klamath River, 2022. Note that RST = rotary screw trap, UPS = upstream, DNS = downstream, and YOY = young-of-the-year.

USFWS 2022 Mainstem Klamath River Outmigrant Trap Juvenile Salmonid Catch Summary

U.S. Fish & Wildlife Service, Arcata Fish & Wildlife Office, 1655 Hendon Road, Arcata, CA 95521, (707)822-7201

Preliminary Data - Subject to Revision

Trap	Calendar week	Sample dates	Q (cfs) <sup>a</sup>		Water temp. (°F) <sup>b</sup>		Trapping days	Chinook ( <i>O. tshawytscha</i> )			Coho ( <i>O. kisutch</i> )			Steelhead ( <i>O. mykiss</i> )			
			Min	Max	Min	Max		YOY			Age 1+			Age 1+			
								No clip	AD clip	Age 1+	YOY	No clip	LM clip	YOY	No clip	AD clip	
Bogus Frame Net	10	3/2-3/4	990	1,020	43.8	44.2	3	74	0	0	0	0	0	0	0	0	0
	11	3/8-3/11	985	996	44.6	45.3	4	120	0	0	0	0	0	0	0	0	0
	12 <sup>c</sup>	3/14-3/18	982	992	-	-	0	-	-	-	-	-	-	-	-	-	-
	13	3/22-3/25	983	995	48.7	51	4	108	0	0	2	0	0	0	0	0	0
	14	3/29-4/1	987	1,190	50.1	50.9	4	483	0	0	44	0	0	0	0	0	0
	15	4/5-4/8	1,290	1,350	50.7	51.6	4	944	0	0	236	0	0	27	0	0	0
	16	4/12-4/15	1,340	1,990	50.9	50.9	1	61	0	0	49	0	0	6	1	0	0
	17 <sup>a</sup>	4/19-4/22	1,650	3,090	-	-	0	-	-	-	-	-	-	-	-	-	-
	18	4/26-4/28	1,310	1,340	53.7	53.9	3	572	0	0	1712	0	0	213	0	0	0
	19	5/3-5/6	1,180	1,180	52.1	53.8	4	347	0	0	445	0	0	549	0	0	0
	20	5/10-5/13	1,180	1,180	51.9	53.7	4	102	0	0	5	0	0	270	0	0	0
	21	5/17-5/20	1,170	1,180	56.1	60	4	65	0	0	21	0	0	370	1	0	0
	22	5/24-5/27	1,170	1,180	55.5	59.3	4	51	25	0	39	0	0	148	0	0	0
23	6/1-6/3	1,020	1,100	59.1	59.3	3	14	1	0	2	0	0	42	0	0	0	
I-5 UPS RST	10	3/1-3/4	964	1,020	41.9	43.8	4	226	0	12	0	0	0	0	1	0	
	11	3/8-3/11	985	996	42.6	44	4	452	0	1	0	0	0	0	0	0	
	12	3/15-3/18	982	990	43.8	44.9	4	291	0	0	0	1	378	0	0	0	
	13	3/22-3/25	983	995	45.2	46.9	4	319	0	5	6	0	2	0	4	0	
	14	3/29-4/1	987	1,190	48.1	49.6	4	416	0	0	19	0	1	0	0	0	
	15	4/5-4/8	1,290	1,350	48.7	50.3	4	212	0	0	12	0	0	1	2	0	
	16	4/12-4/15	1,340	1,990	47.4	49.1	4	831	0	1	8	0	1	5	0	0	
	17	4/19-4/22	1,650	3,090	48.9	49.4	4	1857	0	0	113	1	2	59	1	0	
	18	4/26-4/29	1,250	1,340	52.3	52.8	4	2336	0	0	79	3	1	280	1	0	
	19	5/3-5/6	1,180	1,180	51.2	52.7	4	572	0	0	23	2	4	465	2	0	
	20	5/10-5/13	1,180	1,180	51.4	52.1	4	95	0	0	5	1	2	199	1	0	
	21	5/17-5/20	1,170	1,180	54.3	55.5	4	154	1	0	18	0	3	398	1	0	
	22	5/24-5/27	1,170	1,180	55	57.5	4	345	34	0	36	1	2	280	1	0	
23	6/1-6/3	1,020	1,100	56.3	57.7	3	139	11	0	39	0	0	74	0	0		
I-5 DNS RST	10	3/1-3/4	985	1,020	41.9	43.8	3	104	0	8	0	0	0	0	3	0	
	11	3/8-3/11	964	996	42.6	44	4	207	0	1	0	1	0	0	0	0	
	12	3/15-3/18	982	990	43.8	44.9	3	134	0	0	0	0	123	0	0	0	
	13	3/22-3/25	983	995	45.2	46.9	4	220	0	0	2	0	9	0	3	0	
	14	3/29-4/1	987	1,190	48.1	49.6	4	290	0	0	15	0	2	0	1	0	
	15	4/5-4/8	1,290	1,350	48.7	50.3	4	206	0	0	7	0	0	2	1	0	
	16	4/12-4/15	1,340	1,990	47.4	48.9	4	380	0	0	7	0	1	2	0	0	
	17	4/19-4/22	1,650	3,090	48.9	49.4	4	752	0	0	42	0	0	25	0	0	
	18	4/26-4/29	1,250	1,340	52.3	52.8	4	850	0	1	49	2	0	131	0	0	
	19	5/3-5/6	1,180	1,180	51.2	52.7	4	273	0	0	11	1	0	236	0	0	
	20	5/10-5/13	1,180	1,180	51.4	52.1	4	43	0	0	1	0	0	127	0	0	
	21	5/17-5/20	1,170	1,180	54.3	55.5	4	78	0	0	6	1	0	187	1	0	
	22	5/24-5/27	1,170	1,180	55	57.5	4	160	5	0	21	0	1	153	0	0	
23	6/1-6/3	1,020	1,100	56.3	57.7	2	53	1	0	13	0	0	32	0	0		
I-5 Frame Net	10	3/2-3/4	990	1,020	42.2	43.8	3	23	0	0	0	0	0	0	0	0	
	11	3/8-3/11	985	996	42.6	44	4	103	0	0	0	2	0	0	0	0	
	12	3/15-3/18	982	990	43.8	44.9	4	32	0	0	0	0	7	0	0	0	
	13	3/22-3/25	983	995	45.2	46.9	4	50	0	0	1	0	2	1	0	0	
	14	3/29-4/1	987	1,190	48.1	49.6	4	78	0	0	11	0	0	0	0	0	
	15	4/5-4/8	1,290	1,350	48.7	50.3	2	17	0	0	0	0	0	0	0	0	
	16	4/12-4/15	1,340	1,990	48.9	48.9	1	16	0	0	1	0	0	0	0	0	
	17 <sup>a</sup>	4/19-4/22	1,650	3,090	-	-	0	-	-	-	-	-	-	-	-	-	-
	18	4/26-4/29	1,250	1,340	52.3	52.8	4	1133	0	0	378	0	0	228	0	0	0
	19	5/3-5/6	1,180	1,180	51.2	52.7	4	248	0	0	81	0	0	260	0	0	0
	20	5/10-5/13	1,180	1,180	51.4	52.1	4	22	0	0	3	0	0	130	0	0	0
	21	5/17-5/20	1,170	1,180	54.3	55.5	4	15	0	0	3	0	0	200	0	0	0
	22	5/24-5/27	1,170	1,180	55	57.5	3	28	3	0	9	0	1	121	0	0	0
23	6/1-6/3	1,020	1,100	56.3	57.7	3	19	0	0	1	0	0	11	0	0	0	

<sup>a</sup> mean discharge from day of sampling (discharge below IGD used for Bogus and I-5 sites; flow at Kinsman Site is Klamath River flow at Seiad minus Scott River flow; discharge at Weitchpec Site is discharge near Orleans)

<sup>b</sup> temperature recorded at time of trap check

<sup>c</sup> trap not set this week because trapping operations were limited due to a flow event and/or hatchery release

Table 1 cont. In-season summary of the total catch by week of adipose fin-clipped (AD Clip) and non-adipose fin clipped (No Clip) Chinook Salmon and steelhead and left maxillary-clipped (LM Clip) and non-maxillary clipped (No Clip) Coho Salmon by trap at the Bogus, I 5, and Kinsman trap sites on the mainstem Klamath River, 2022. Note that RST = rotary screw trap, UPS = upstream, DNS = downstream, and YOY = young-of-the-year.

USFWS 2022 Mainstem Klamath River Outmigrant Trap Juvenile Salmonid Catch Summary (continued)

U.S. Fish & Wildlife Service, Arcata Fish & Wildlife Office, 1655 Heindon Road, Arcata, CA 95521, (707)822-7201

Preliminary Data - Subject to Revision

Trap	Calendar week	Sample dates	Q (cfs) <sup>a</sup>		Water temp. (F) <sup>b</sup>		Trapping days	Chinook ( <i>O. tshawytscha</i> )			Coho ( <i>O. kisutch</i> )			Steelhead ( <i>O. mykiss</i> )		
			Min	Max	Min	Max		YOY			YOY	Age 1 +		YOY	Age 1 +	
								No clip	AD clip	Age 1+		No clip	LM clip		No clip	AD clip
Kinsman RST	10	3/1-3/4	1,338	1,432	45.0	47.2	4	162	0	1	3	8	0	5	3	0
	11	3/8-3/11	1,330	1,349	43.3	46.5	4	144	0	0	2	5	0	1	0	0
	12	3/15-3/18	1,416	1,438	44.7	47.8	4	109	0	0	1	5	6	0	5	0
	13	3/22-3/25	1,416	1,503	47.1	52.8	4	191	0	0	5	2	45	4	7	0
	14	3/29-4/1	1,449	1,498	49.1	51.4	4	349	0	0	37	4	2	0	11	0
	15	4/5-4/8	1,733	1,791	49.2	55.5	4	325	0	0	4	7	2	2	3	0
	16	4/12-4/15	1,750	1,804	47.4	49.8	4	179	0	2	2	2	1	2	2	0
	17*	4/19-4/22	2,654	4,004	-	-	0	-	-	-	-	-	-	-	-	-
	18	4/26-4/29	1,913	2,015	52.1	58.4	4	148	0	2	4	1	5	4	8	0
	19	5/3-5/6	1,782	2,030	56.9	60.1	4	123	0	1	46	6	4	27	14	0
	20	5/10-5/13	1,838	1,955	50.1	55.2	4	77	0	1	14	3	5	20	7	0
	21	5/17-5/20	1,875	1,930	57.7	60.9	4	51	0	2	44	2	1	9	2	0
	22	5/24-5/27	1,815	1,876	61.7	61.8	4	65	0	0	20	1	2	11	7	0
23	6/1-6/3	1,569	1,710	61.4	63.8	2	31	0	0	36	0	1	35	6	0	
Weitchpec RST	10	3/1-3/4	4,194	4,597	-	-	4	265	0	2	0	1	0	0	5	0
	11	3/8-3/11	3,461	3,687	46.2	46.6	4	203	0	0	2	0	0	0	2	0
	12	3/15-3/18	4,470	5,610	48.2	48.4	1	18	0	0	0	0	0	0	6	0
	13	3/22-3/24	4,110	5,290	48.9	50.5	3	138	0	0	0	1	16	0	15	0
	14	3/29-4/1	4,480	5,190	50.2	52.0	4	84	0	0	0	1	0	0	20	0
	15	4/5-4/8	4,730	6,000	48.2	51.6	4	82	0	0	0	0	2	2	39	0
	16	4/12-4/15	4,560	5,280	49.5	49.6	2	67	0	0	0	1	0	0	11	0
	17*	4/19-4/22	8,750	11,600	-	-	0	-	-	-	-	-	-	-	-	-
	18	4/26-4/29	7,380	8,720	50.0	51.8	4	27	0	0	4	1	0	0	3	0
	19	5/3-5/6	7,260	11,400	49.8	50.5	3	17	0	0	1	2	3	2	12	0
	20	5/10-5/13	4,620	8,280	49.1	49.6	3	24	0	0	0	2	0	0	5	0
	21	5/17-5/20	7,160	8,060	52.7	54.5	4	48	0	7	0	8	6	8	11	0
	22	5/24-5/27	6,240	6,470	56.3	58.1	4	151	0	31	3	13	1	5	37	0
23	6/1-6/3	4,780	5,070	58.1	59.0	3	13	0	83	4	15	2	4	106	0	
Weitchpec US Frame	11	3/8-3/11	3,461	3,687	46.2	46.6	4	94	0	0	0	0	0	0	0	0
	12	3/15-3/18	4,470	5,610	-	-	0	-	-	-	-	-	-	-	-	-
	13	3/22-3/24	4,110	5,290	48.9	50.5	3	107	0	0	0	0	0	0	0	0
	14	3/30-4/1	4,480	5,130	50.2	50.5	3	50	0	0	0	0	0	0	0	0
	15	4/7-4/8	4,730	4,940	50.5	51.6	2	30	0	0	0	0	0	0	0	0
	16	4/12-4/15	4,560	5,280	49.5	49.6	2	41	0	0	0	0	0	0	0	0
	17*	4/19-4/22	8,750	11,600	-	-	0	-	-	-	-	-	-	-	-	-
	18*	4/26-4/29	7,380	8,720	-	-	0	-	-	-	-	-	-	-	-	-
	19*	5/3-5/6	7,260	11,400	-	-	0	-	-	-	-	-	-	-	-	-
	20*	5/10-5/13	4,620	8,280	-	-	0	-	-	-	-	-	-	-	-	-
	21	5/17-5/20	7,160	8,060	-	-	0	-	-	-	-	-	-	-	-	-
	22	5/24-5/27	6,240	6,470	-	-	0	-	-	-	-	-	-	-	-	-
	23	6/2-6/3	4,780	4,960	59.0	59.0	2	3	0	0	1	0	0	5	0	0

<sup>a</sup> mean discharge from day of sampling (discharge below IGD used for Bogus and I-5 sites; flow at Kinsman Site is Klamath River flow at Seiad minus Scott River flow; discharge at Weitchpec Site is discharge near Orleans)

<sup>b</sup> temperature recorded at time of trap check

<sup>c</sup> trap not set this week because trapping operations were limited due to a flow event and/or hatchery release



Table 2 cont. In-season summary of the average catch-per-day by week of non-adipose fin-clipped (No Clip) and adipose fin-clipped (AD Clip) Chinook Salmon and steelhead and non-maxillary clipped (No Clip) and left maxillary-clipped (LM Clip) Coho Salmon by trap at the Bogus, I-5, and Kinsman trap sites on the mainstem Klamath River, 2022. Note that RST = rotary screw trap, UPS = upstream, DNS = downstream, and YOY = young-of-the-year.

USFWS 2022 Mainstem Klamath River Outmigrant Trap Juvenile Salmonid Catch-per-Day Summary (continued)

U.S. Fish & Wildlife Service, Arcata Fish & Wildlife Office, 1655 Heindon Road, Arcata, CA 95521, (707)822-7201

Preliminary Data - Subject to Revision

Trap	Calendar week	Sample dates	Q (cfs) <sup>a</sup>		Water temp. (F) <sup>b</sup>		Trapping days	Chinook ( <i>O. tshawytscha</i> )			Coho ( <i>O. kisutch</i> )			Steelhead ( <i>O. mykiss</i> )				
			Min	Max	Min	Max		YOY			Age 1+			Age 1+				
								No clip	AD clip	Age 1+	YOY	No clip	LM clip	YOY	No clip	AD clip		
Kinsman RST	10	3/1-3/4	1,338	1,432	45.0	47.2	4	40.50	0.00	0.25	0.75	2.00	0.00	1.25	0.75	0.00		
	11	3/8-3/11	1,410	1,349	43.3	46.5	4	36.00	0.00	0.00	0.50	1.25	0.00	0.25	0.00	0.00		
	12	3/15-3/18	1,416	1,438	44.7	47.8	4	27.25	0.00	0.00	0.25	1.25	1.50	0.00	1.25	0.00		
	13	3/22-3/25	1,416	1,503	47.1	52.8	4	47.75	0.00	0.00	1.25	0.50	11.25	1.00	1.75	0.00		
	14	3/29-4/1	1,449	1,498	49.1	51.4	4	87.25	0.00	0.00	9.25	1.00	0.50	0.00	2.75	0.00		
	15	4/5-4/8	1,733	1,791	49.2	55.5	4	81.25	0.00	0.00	1.00	1.75	0.50	0.50	0.75	0.00		
	16	4/12-4/15	1,750	1,804	47.4	49.8	4	44.75	0.00	0.50	0.50	0.50	0.25	0.50	0.50	0.00		
	17 <sup>c</sup>	4/19-4/22	2,654	4,004	-	-	0	-	-	-	-	-	-	-	-	-	-	
	18	4/26-4/29	1,913	2,015	52.1	58.4	4	37.00	0.00	0.50	1.00	0.25	1.25	1.00	2.00	0.00		
	19	5/3-5/6	1,782	2,030	56.9	60.1	4	30.75	0.00	0.25	11.50	1.50	1.00	6.75	3.50	0.00		
	20	5/10-5/13	1,838	1,955	50.1	55.2	4	19.25	0.00	0.25	3.50	0.75	1.25	5.00	1.75	0.00		
	21	5/17-5/20	1,875	1,930	57.7	60.9	4	12.75	0.00	0.50	11.00	0.50	0.25	2.25	0.50	0.00		
	22	5/24-5/27	1,815	1,876	61.7	61.8	4	16.25	0.00	0.00	5.00	0.25	0.50	2.75	1.75	0.00		
	23	6/1-6/3	1,569	1,710	61.4	63.8	2	15.50	0.00	0.00	18.00	0.00	0.50	17.50	3.00	0.00		
	Weitchpec RST	10	3/1-3/4	4,194	4,597	-	-	4	66.25	0.00	0.50	0.00	0.25	0.00	0.00	1.25	0.00	
		11	3/8-3/11	3,461	3,687	46.2	46.6	4	50.75	0.00	0.50	0.00	0.25	0.00	0.00	0.50	0.00	
		12	3/15-3/18	4,470	5,610	48.2	48.4	1	18.00	0.00	0.00	0.00	0.00	0.00	0.00	6.00	0.00	
		13	3/22-3/24	4,110	5,290	48.9	50.5	3	46.00	0.00	0.00	0.00	0.33	5.33	0.00	5.00	0.00	
		14	3/29-4/1	4,480	5,190	50.2	52.0	4	21.00	0.00	0.00	0.00	0.25	0.00	0.00	5.00	0.00	
		15	4/5-4/8	4,730	6,000	48.2	51.6	4	20.50	0.00	0.00	0.00	0.00	0.50	0.50	9.75	0.00	
		16	4/12-4/15	4,560	5,280	49.5	49.6	2	33.50	0.00	0.00	0.00	0.50	0.00	0.00	5.50	0.00	
		17 <sup>c</sup>	4/19-4/22	8,750	11,600	-	-	0	-	-	-	-	-	-	-	-	-	-
		18	4/26-4/29	7,380	8,720	50.0	51.8	4	6.75	0.00	0.00	1.00	0.25	0.00	0.00	0.75	0.00	
19		5/3-5/6	7,260	11,400	49.8	50.5	3	5.67	0.00	0.00	0.33	0.67	1.00	0.67	4.00	0.00		
20		5/10-5/13	4,620	8,280	49.1	49.6	3	8.00	0.00	0.00	0.00	0.67	0.00	0.00	1.67	0.00		
21		5/17-5/20	7,160	8,060	52.7	54.5	4	12.00	0.00	1.75	0.00	2.00	1.50	2.00	2.75	0.00		
22		5/24-5/27	6,240	6,470	56.3	58.1	4	37.75	0.00	7.75	0.75	3.25	0.25	1.25	9.25	0.00		
23		6/1-6/3	4,780	5,070	58.1	59.0	3	4.33	0.00	27.67	1.33	5.00	0.67	1.33	35.33	0.00		
Weitchpec US Frame		11	3/8-3/11	3,461	3,687	46.2	46.6	4	23.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
		12	3/15-3/18	4,470	5,610	-	-	0	-	-	-	-	-	-	-	-	-	
		13	3/22-3/24	4,110	5,290	48.9	50.5	3	35.67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
		14	3/30-4/1	4,480	5,130	50.2	50.5	3	16.67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
		15	4/7-4/8	4,730	4,940	50.5	51.6	2	15.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
		16	4/12-4/15	4,560	5,280	49.5	49.6	2	20.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
		17 <sup>c</sup>	4/19-4/22	8,750	11,600	-	-	0	-	-	-	-	-	-	-	-	-	
		18 <sup>c</sup>	4/26-4/29	7,380	8,720	-	-	0	-	-	-	-	-	-	-	-	-	
		19 <sup>c</sup>	5/3-5/6	7,260	11,400	-	-	0	-	-	-	-	-	-	-	-	-	
	20	5/10-5/13	4,620	8,280	-	-	0	-	-	-	-	-	-	-	-	-		
	21	5/17-5/20	7,160	8,060	-	-	0	-	-	-	-	-	-	-	-	-		
	22	5/24-5/27	6,240	6,470	-	-	0	-	-	-	-	-	-	-	-	-		
23	6/2-6/3	4,780	4,960	59.0	59.0	2	1.50	0.00	0.00	0.50	0.00	0.00	2.50	0.00	0.00			

<sup>a</sup> mean daily discharge range during sampling dates (discharge below IGD used for Bogus and I-5 sites; flow at Kinsman Site is Klamath River flow at Seiad minus Scott River flow; discharge at Weitchpec Site is discharge near Orleans)

<sup>b</sup> temperature recorded at time of trap check

<sup>c</sup> trap not set this week because trapping operations were limited due to a flow event and/or hatchery release



Table 4. In-season summary of clinical signs of disease in young-of-the-year Chinook Salmon by site at the Bogus, I-5, and Kinsman sites on the mainstem Klamath River, 2022. *Note: Although only Chinook Salmon are reported in this table, we also monitor clinical signs of diseases in Coho Salmon and other species.*

**USFWS 2022 Mainstem Klamath River YOY Chinook Salmon Clinical Signs of Disease Summary**

U.S. Fish & Wildlife Service, Arcata Fish & Wildlife Office, 1655 Heindon Road, Arcata, CA 95521, (707)822-7201

Preliminary Data - Subject to Revision

Site	Calendar week	Sampling dates	Weekly mean flow (cfs) <sup>a</sup>	Water temp. (°F) <sup>b</sup>		Belly condition			Gills		Condition			
				Min	Max	Sample size	Distended		Sample size	Color		Eroded or fungal		
							# positive	%		# positive	%	# positive	%	
Bogus	10	3/2-3/4	980	43.8	44.2	40	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	11	3/8-3/11	994	44.6	45.3	51	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	12 <sup>d</sup>	3/14-3/18	993	-	-	-	-	-	-	-	-	-	-	
	13	3/22-3/25	995	48.7	51.0	70	0	0.0%	1	0	- <sup>c</sup>	0	- <sup>c</sup>	
	14	3/29-4/1	1,061	50.1	50.9	87	0	0.0%	2	0	- <sup>c</sup>	0	- <sup>c</sup>	
	15	4/5-4/8	1,310	50.7	51.6	90	0	0.0%	7	0	- <sup>c</sup>	0	- <sup>c</sup>	
	16	4/12-4/15	1,877	50.9	50.9	30	0	0.0%	5	0	- <sup>c</sup>	0	- <sup>c</sup>	
	17 <sup>d</sup>	4/19-4/22	2,599	-	-	-	-	-	-	-	-	-	-	
	18	4/26-4/28	1,309	53.7	53.9	89	8	9.0%	23	1	- <sup>c</sup>	0	- <sup>c</sup>	
	19	5/3-5/6	1,211	52.1	53.8	89	1	1.1%	30	0	0.0%	2	6.7%	
	20	5/10-5/13	1,179	51.9	53.7	127	6	4.7%	55	1	1.8%	1	1.8%	
	21	5/17-5/20	1,176	56.1	60.0	40	16	40.0%	36	1	2.8%	1	2.8%	
	22	5/24-5/27	1,176	55.5	59.3	58	5	8.6%	45	0	0.0%	0	0.0%	
	23	6/1-6/3	1,100	59.1	59.3	8	1	- <sup>c</sup>	5	0	- <sup>c</sup>	0	- <sup>c</sup>	
	I-5	10	3/1-3/4	980	41.9	43.8	98	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>
		11	3/8-3/11	994	42.6	44.0	161	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>
		12	3/15-3/18	993	43.8	44.9	114	0	0.0%	2	0	- <sup>c</sup>	0	- <sup>c</sup>
		13	3/22-3/25	995	45.2	46.9	128	0	0.0%	2	0	- <sup>c</sup>	0	- <sup>c</sup>
		14	3/29-4/1	1,061	48.1	49.6	158	0	0.0%	7	0	- <sup>c</sup>	0	- <sup>c</sup>
		15	4/5-4/8	1,310	48.7	50.3	93	0	0.0%	36	0	0.0%	0	0.0%
		16	4/12-4/15	1,877	47.4	49.1	151	1	0.7%	111	1	0.9%	0	0.0%
		17	4/19-4/22	2,599	48.9	49.4	87	0	0.0%	68	0	0.0%	0	0.0%
		18	4/26-4/29	1,309	52.3	52.8	177	1	0.6%	110	0	0.0%	0	0.0%
19		5/3-5/6	1,211	51.2	52.7	160	0	0.0%	110	0	0.0%	0	0.0%	
20		5/10-5/13	1,179	51.4	52.1	96	6	6.3%	85	7	8.2%	8	9.4%	
21		5/17-5/20	1,176	54.3	55.5	82	4	4.9%	67	2	3.0%	3	4.5%	
22		5/24-5/27	1,176	55.0	57.5	68	8	11.8%	60	4	6.7%	2	3.3%	
23	6/1-6/3	1,100	56.3	57.7	58	5	8.6%	54	5	9.3%	5	9.3%		
Kinsman	10	3/1-3/4	1,367	45.0	47.2	85	0	0.0%	3	0	- <sup>c</sup>	0	- <sup>c</sup>	
	11	3/8-3/11	1,345	43.3	46.5	85	0	0.0%	12	0	- <sup>c</sup>	0	- <sup>c</sup>	
	12	3/15-3/18	1,410	44.7	47.8	67	0	0.0%	13	0	- <sup>c</sup>	0	- <sup>c</sup>	
	13	3/22-3/25	1,444	47.1	52.8	90	0	0.0%	41	0	0.0%	0	0.0%	
	14	3/29-4/1	1,507	49.1	51.4	90	0	0.0%	66	0	0.0%	0	0.0%	
	15	4/5-4/8	1,752	49.2	55.5	90	0	0.0%	78	0	0.0%	0	0.0%	
	16	4/12-4/15	2,070	47.4	49.8	90	1	1.1%	88	0	0.0%	0	0.0%	
	17 <sup>d</sup>	4/19-4/22	3,311	-	-	-	-	-	-	-	-	-	-	
	18	4/26-4/29	1,978	52.1	58.4	85	0	0.0%	82	0	0.0%	0	0.0%	
	19	5/3-5/6	1,939	56.9	60.1	74	3	4.1%	69	0	0.0%	0	0.0%	
	20	5/10-5/13	1,930	50.1	55.2	59	0	0.0%	59	1	1.7%	0	0.0%	
	21	5/17-5/20	1,888	57.7	60.9	32	5	15.6%	31	1	3.2%	0	0.0%	
	22	5/24-5/27	1,838	61.7	61.8	33	4	12.1%	33	0	0.0%	0	0.0%	
23	6/1-6/3	1,709	61.4	63.8	20	2	- <sup>c</sup>	19	2	- <sup>c</sup>	1	- <sup>c</sup>		
Weitchpec	10	3/1-3/4	4,167	-	-	120	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	11	3/8-3/11	3,644	46.2	46.6	195	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	12	3/15-3/18	4,542	48.2	48.4	7	0	- <sup>c</sup>	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	13	3/22-3/24	4,750	48.9	50.5	176	0	0.0%	0	-	- <sup>c</sup>	-	- <sup>c</sup>	
	14	3/29-4/1	4,937	50.2	52.0	105	0	0.0%	9	0	- <sup>c</sup>	0	- <sup>c</sup>	
	15	4/5-4/8	5,024	48.2	51.6	59	0	0.0%	26	0	- <sup>c</sup>	0	- <sup>c</sup>	
	16	4/12-4/15	4,914	49.5	49.6	40	0	0.0%	32	0	0.0%	0	0.0%	
	17 <sup>d</sup>	4/19-4/22	9,094	-	-	-	-	-	-	-	-	-	-	
	18	4/26-4/29	8,243	50.0	51.8	13	0	- <sup>c</sup>	2	0	- <sup>c</sup>	0	- <sup>c</sup>	
	19	5/3-5/6	9,256	49.8	50.5	16	0	- <sup>c</sup>	7	0	- <sup>c</sup>	0	- <sup>c</sup>	
	20	5/10-5/13	8,013	49.1	49.6	20	0	- <sup>c</sup>	20	0	- <sup>c</sup>	0	- <sup>c</sup>	
	21	5/17-5/20	6,353	52.7	54.5	20	0	- <sup>c</sup>	20	0	- <sup>c</sup>	0	- <sup>c</sup>	
	22	5/24-5/27	5,266	56.3	58.1	40	2	5.0%	40	0	0.0%	0	0.0%	
23	6/1-6/3	7,650	58.1	59.0	46	0	0.0%	46	0	0.0%	0	0.0%		

<sup>a</sup> discharge below IGD used for Bogus and I-5 sites; discharge at Kinsman Site is Klamath River discharge near Seiad Valley minus discharge in the Scott River near Fort Jones; discharge at Weitchpec Site is discharge near Orleans

<sup>b</sup> temperature recorded at time of trap check/seine

<sup>c</sup> sample size too low for a reportable calculation

<sup>d</sup> trap not set this week because trapping operations were limited due to a flow event and/or hatchery release






Figure 1. Daily mean discharge below Iron Gate Dam, Klamath River (USGS Gaging Station 11516530) from late February through July, 2000–2022.




Figure 2. Klamath River daily mean discharge at the Kinsman Trap Site from late February through July 2000–2020. Flow measurements are not available at this location. Therefore, Klamath River flow near Seiad Valley, California (USGS Gaging Station 11520500) minus flow from the Scott River near Fort Jones, California (USGS 11519500) is used as a surrogate.




Figure 3. Daily mean discharge of Klamath River at Orleans, California (USGS Gaging Station 10523000) from late February through July 2000–2022.