

**Recovery Plan for *Eriogonum ovalifolium* var. *williamsiae* (Steamboat buckwheat)**  
[https://ecos.fws.gov/docs/recovery\\_plan/950920b.pdf](https://ecos.fws.gov/docs/recovery_plan/950920b.pdf)

**Original Approved:** September 29, 1995  
**Original Prepared by:** Nevada State Office

**DRAFT AMENDMENT**

We have identified best available information that indicates the need to amend recovery criteria for *Eriogonum ovalifolium* var. *williamsiae* (Steamboat buckwheat) since the recovery plan was completed. In this proposed modification, we synthesize the adequacy of the existing recovery criteria, show amended recovery criteria, the rationale supporting the proposed recovery plan modification, and analyses of the most recent monitoring data. The proposed modification is shown as an addendum that supplements the recovery plan, superseding only p. iii of the recovery plan.

**For**  
**U.S. Fish and Wildlife Service**  
**Pacific Southwest Region**  
**Reno, Nevada**  
  
**December 2018**

Approved: XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

Regional Director, Pacific Southwest Region, Region 8,  
U.S. Fish and Wildlife Service

Date: XXXXXXXXXXXXXXXXXXXX

## METHODOLOGY USED TO COMPLETE THE RECOVERY PLAN AMENDMENT

This amendment was prepared by the Reno Fish and Wildlife Office. We used information in our files, data from the Bureau of Land Management (BLM), the Nevada geospatial database maintained by Nevada Natural Heritage Program, and information from species experts. The amended criteria will be peer reviewed in accordance with the OMB Peer Review Bulletin following the publication of the Notice of Availability.

## ADEQUACY OF RECOVERY CRITERIA

Section 4(f)(1)(B)(ii) of the Endangered Species Act (Act) requires that each recovery plan shall incorporate, to the maximum extent practicable, “objective, measurable criteria which, when met, would result in a determination...that the species be removed from the list.” Legal challenges to recovery plans (see *Fund for Animals v. Babbitt*, 903 F. Supp. 96 (D.D.C. 1995)) and a Government Accountability Audit (GAO 2006) also have affirmed the need to frame recovery criteria in terms of threats assessed under the five factors.

### Recovery Criteria

See previous version of criteria in recovery plan for *Eriogonum ovalifolium* var. *williamsiae* on p. iii, in the recovery plan, available [here](#).

### Synthesis

*Eriogonum ovalifolium* var. *williamsiae* is a low, densely matted, compact perennial plant in the buckwheat family (Polygonaceae). This species is known from a single population in Washoe County, Nevada located approximately 10 miles south of Reno. *Eriogonum ovalifolium* var. *williamsiae* is endemic to substrates derived from hot springs deposits known as sinter in the Steamboat Hills, which is an area of significant geothermal activity that once featured the largest concentration of geysers in the United States outside of Yellowstone National Park (BLM 1993). The species occupies a total of approximately 50 acres in an area of approximately 250-370 acres on lands that are managed by the Bureau of Land Management, State of Nevada, and private owners (Service 2009). About half of the lands occupied by *E. ovalifolium* var. *williamsiae* are private and under a 30-year lease by Ormat Technologies Inc. for geothermal power production. The remainder of its habitat occurs on lands under management by the BLM including the Steamboat Hot Springs Geyser Basin Area of Critical Environmental Concern (ACEC; Service 1995), proposed to be changed to the Steamboat Buckwheat Botanical ACEC in the Carson City BLM Resource Management Plan Revision (BLM 2014).

The overall distribution of *E. ovalifolium* var. *williamsiae* has changed little since the time of listing in 1986. At the time of our last 5-Year Review, the recovery priority number was raised from 6c to 3 because threats to the species remain high, but biological and ecological needs of *Eriogonum ovalifolium* var. *williamsiae* are better understood than they were at the time of listing (Service 2009). One of the threats identified in the original listing rule has not materialized (i.e. development of a park on a BLM parcel leased to the Washoe County Parks and Recreation Department), while others (i.e. off-road vehicle activity, refuse dumping, and potential mining) have had less dire consequences than predicted due to fencing of the geothermal site (Service 2009). Other threats identified at the time of listing – drilling of

geothermal test wells, commercial development on private lands adjacent to a colony of plants, and changes in moisture availability continue to be of concern. In 2011, an eleven acre private parcel was transferred to the State of Nevada, of which 6.7 acres are occupied by *E. ovalifolium* var. *williamsiae* (C. Clark, unpublished data 2004, Washoe County 2018). In 2017, BLM parcels occupied by *E. ovalifolium* var. *williamsiae* were leased with a no surface occupancy stipulation for geothermal resources, however, there is potential for geothermal exploration and/or development activities to indirectly impact the species through surface water flows and altering soil conditions (BLM 2017, BLM 2018).

Various estimates of the abundance of *Eriogonum ovalifolium* var. *williamsiae* have been made over the 32 years since the plant was listed. These estimates have ranged from 10,000 to 15,000 individual plants in the final rule listing the species (Service 1986) to 85,000 plants by Knight (1997), to 200,000 plants by Morefield (2001). Knight 1993 noted that a precise enumeration of individual plants is infeasible because the species is rhizomatous and propagates primarily by vegetative runners.

In 2003, a demographic monitoring program was implemented with sampling conducted along permanent transects in two areas within the population – one in the Central Drainage and the other on the Main Terrace (Pavlik 2002, Pavlik and Stanton 2003). Data were collected in May 2003, June 2005, and July 2017. On the Central Drainage, all size classes of *Eriogonum ovalifolium* var. *williamsiae* have been declining since 2003, especially smaller plants. On the Main Terrace, both small and medium sized plants have been declining, while larger plants have been increasing since baseline measurements in 2003. Additionally, though not a threat at the time of listing, nonnative weed species have started to colonize the sinter substrate. Though weed cover is low, *Bromus tectorum* (cheatgrass) on the Central Drainage and *Vulpia myuros* (rattail fescue) on the Main Terrace has increased over time. Cover of trees, shrubs, and perennial grasses have remained unchanged over the 14 year period of monitoring data, suggesting that colonization by other native species in this habitat is not occurring. Further monitoring results are discussed in Appendix A.

In 1993, the Nature Conservancy developed the *Steamboat Buckwheat Management Plan* (Knight 1993). However, the plan is limited in scope to the private land under lease for the geothermal power production and limited in time from 1991–2021. In 2005, Pavlik *et al.* developed *A Management Plan for the Steamboat Buckwheat* that covers all lands occupied by *Eriogonum ovalifolium* var. *williamsiae* and offered a vision of restoration of the Steamboat Springs geothermal ecosystem for public enjoyment and education through restoration of ecosystem integrity in a way that sustains power production. However, implementation of the plan in full has not been achieved. Some aspects of the plan (i.e. public education and access opportunities) are unlikely to be implemented because of safety concerns with allowing public access by the geothermal company (Service 2009).

## **AMENDED RECOVERY CRITERIA**

Recovery criteria serve as objective, measurable guidelines to assist in determining when an endangered species has recovered to the point that it may be downlisted to threatened, or that the protections afforded by the Act are no longer necessary and *Eriogonum ovalifolium* var.

*williamsiae* may be delisted. Delisting is the removal of a species from the Federal Lists of Endangered and Threatened Wildlife and Plants. Downlisting is the reclassification of a species from an endangered species to a threatened species. The term “endangered species” means any species (species, sub-species, or distinct population segment (DPS)) which is in danger of extinction throughout all or a significant portion of its range. The term “threatened species” means any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.

Revisions to the Lists, including delisting or downlisting a species, must reflect determinations made in accordance with sections 4(a)(1) and 4(b) of the Act. Section 4(a)(1) requires that the Secretary determine whether a species is an endangered species or threatened species (or not) because of threats to the species. Section 4(b) of the Act requires that the determination be made “solely on the basis of the best scientific and commercial data available.” Thus, while recovery plans provide important guidance to the Service, States, and other partners on methods of minimizing threats to listed species and measurable objectives against which to measure progress towards recovery, they are guidance and not regulatory documents.

Recovery criteria should help indicate when we would anticipate that an analysis of the species’ status under section 4(a)(1) would result in a determination that the species is no longer an endangered species or threatened species. A decision to revise the status of or remove a species from the Federal Lists of Endangered and Threatened Wildlife and Plants, however, is ultimately based on an analysis of the best scientific and commercial data then available, regardless of whether that information differs from the recovery plan, which triggers rulemaking. When changing the status of a species, we first propose the action in the *Federal Register* to seek public comment and peer review, followed by a final decision announced in the *Federal Register*.

Downlisting criteria were included in the Steamboat Buckwheat Recovery Plan (Service 1995). We provide delisting criteria for *Eriogonum ovalifolium* var. *williamsiae*, which were not included in the Steamboat Buckwheat Recovery Plan, as follows:

#### **Current Downlisting Recovery Criteria (from original recovery plan)**

Downlisting for *Eriogonum ovalifolium* var. *williamsiae* can be considered when:

1. Protective conservation easements or fee acquisitions secure approximately 185 acres of occupied habitat currently in private ownership;
2. Cooperative agreements are established for approximately 80 acres of occupied public lands and approximately 37 acres of occupied State lands within a highway easement; and
3. Comprehensive management plans have been developed and implemented on all occupied habitat.

#### **Delisting Recovery Criteria (amended recovery criteria)**

Delisting may be warranted when the current downlisting criteria have been met and the species exhibits sufficient resiliency, redundancy, and representation to support long-term viability.

When all downlisting criteria have been met for *Eriogonum ovalifolium* var. *williamsiae*, it can be considered for delisting if all three of the additional measures described below are attained:

1. Threats are reduced or eliminated so that the species is capable of persisting without substantial human intervention or perpetual endowments are secured for management necessary to maintain the continued existence of the species. Outstanding management needs include: a) implementing the monitoring protocol, b) updating and renewing the *Steamboat Buckwheat Management Plan* (Knight 1997), c) controlling competition with nonnative weeds, , and d) exploring potential methods of restoration of geothermal processes that maintain and create habitat.
2. All size classes are represented, the population is increasing or stable, and management objectives identified in the demographic monitoring protocol are achieved (Pavlik 2002, Pavlik and Stanton 2003). Monitoring objectives were developed to detect and document 1) trends in the numbers of *Eriogonum ovalifolium* var. *williamsiae* plants in typical habitats, 2) the frequency and contribution of episodic reproduction to population stability, and 3) successional changes in common species that comprise the plant community of Steamboat Hills. Monitoring objectives are as follows:
  - a. *Eriogonum ovalifolium* var. *williamsiae* in the Main Terrace and Central Drainage habitats are within  $\pm 15\%$  of their 2003 baseline levels after five and ten consecutive years of monitoring.
  - b. *Eriogonum ovalifolium* var. *williamsiae* in the Main Terrace and Central Drainage habitats each produce a significant cohort of seedlings at least once during five consecutive years of monitoring (or twice in ten years).
  - c. Total live absolute cover by subpopulations of common shrubs [e.g. *Artemisia tridentata* (sagebrush), *Atriplex confertifolia* (shadscale), *Chrysothamnus nauseosus* (rabbitbrush), *Purshia tridentata* (bitterbrush)], perennial grasses [e.g. *Poa secunda* (bluegrass), *Leymus cinereus* (Great Basin wildrye)], and weeds [e.g. *Bromus tectorum* (cheatgrass)], as well as *E. ovalifolium* var. *williamsiae* are within  $\pm 15\%$  of their 2003 baseline levels after five and ten consecutive years of monitoring.
3. The *ex situ* seedbank is maintained through the collection of fresh seed from *Eriogonum ovalifolium* var. *williamsiae* plants every 10 years. Collections that are spread over time produce lower extinction risk to wild populations, while maintaining a species genetic variation within an *ex situ* seedbank (Menges *et al.* 2004). The *ex situ* seedbank is currently maintained with Center for Plant Conservation-affiliated botanic garden, the Rae Selling Berry Seed Bank and Plant Conservation Program at Portland State University (formally the Berry Botanic Garden). Currently, the *ex situ* seedbank holds approximately 23,000 viable seeds from collections made from 1992-1995 in long-term storage (i.e. freezer). In 1999, a germination trial on a sample of these seeds found greater than 70% germination (E. Guerrant, Rae Selling Berry Seed Bank, pers. comm. 2018a, E. Guerrant, unpubl. data, 2018b).

### **Rationale for Amended Recovery Criteria**

We have amended the recovery criteria for *Eriogonum ovalifolium* var. *williamsiae* to include delisting criteria that incorporates the biodiversity principles of representation, resiliency, and

redundancy (Schaffer and Stein 2000) and threats addressed under the five factors. The amended criteria were developed based on the Service's current understanding of the species needs and requirements. This understanding includes information gathered since the original recovery plan was published, such as recent population monitoring and trends, along with an updated understanding of threats affecting this species. The criteria presented are based on an improved understanding of biological and ecological factors and the reduction of threats to the species; they include a temporal aspect to ensure that the species is resilient to expected variation within a reasonable time frame.

## LITERATURE CITED

- [BLM] Bureau of Land Management. 1993. Steamboat Hot Springs Geyser Area of Critical Environmental Concern and recreation activity management plan. Carson City Field Office, Carson City, Nevada. 15 pp.
- [BLM] Bureau of Land Management. 2014. Draft Resource Management Plan and Environmental Impact Statement. Volume 2: Chapter 4. Carson City District Office, Carson City, Nevada. November 2014. 828 pp.
- [BLM] Bureau of Land Management. 2017. Decision Record: Geothermal Leasing, Washoe County Parcels October 2016 Lease Sales EA. Carson City Field Office, Carson City District Office, Carson City, Nevada. August 2017. 28 pp.
- [BLM] Bureau of Land Management. 2018. Land & Mineral System Reports [web application]. Washington, DC. Available online at < <https://reports.blm.gov/reports/LR2000/>> Accessed 24 September 2018.
- Clark, C. 2004. JBR Environmental Consultants, Inc., Reno, Nevada. Unpublished data. 3 pp.
- Guerrant, E. 2018a. Director, Rae Selling Berry Seed Bank, Portland State University, Portland, Oregon. Phone call with Sarah Kulpa, Reno Fish and Wildlife Office, Reno, Nevada, on September 25, 2018. Subject: *Eriogonum ovalifolium* var. *williamsiae* ex situ seedbank.
- Guerrant, E. 2018b. Rae Selling Berry Seed Bank, Portland State University, Portland Oregon. Unpublished data.
- Knight, T.A. 1997. Steamboat buckwheat management plan. Report submitted to SB Geo, Inc., Reno, Nevada. 8 pp.
- Menges, E.S., E.O. Guerrant Jr., and S. Hamzé. Effects of seed collection on the extinction risk of perennial plants. Pages 305-324 in E.O. Guerrant Jr., K. Havens, and M. Maunder (editors), Ex situ plant conservation: supporting species survival in the wild. Island Press, Washington, DC.
- Morefield, J.D. (editor) 2001. Rare Plant Fact Sheet – *Eriogonum ovalifolium* var. *williamsiae* in *Nevada Rare Plant Atlas*. Nevada Natural Heritage Program, Carson City, Nevada compiled for U.S. Fish and Wildlife Service, Portland, Oregon and Reno, Nevada.
- Pavlik, B.M. 2002. A monitoring program form the Steamboat buckwheat (*Eriogonum ovalifolium* var. *williamsiae*), Washoe County, Nevada. Unpublished report prepared for the United States Fish and Wildlife Service, Reno, Nevada. 36 pp.

- Pavlik, B.M. and A.E. Stanton. 2003. Implementation of a monitoring program for the Steamboat buckwheat (*Eriogonum ovalifolium* var. *williamsiae*), Washoe County, Nevada. I. Establishing Baseline Year One. Unpublished report prepared for Nevada Division of Forestry, Carson City, Nevada, and Nevada Fish and Wildlife Office, Reno, Nevada. 81 pp.
- Pavlik, B.M., A.E. Stanton, and M.J. Bernegger. 2005. A management plan for the steamboat buckwheat (*Eriogonum ovalifolium* var. *williamsiae*) Washoe County, Nevada. Unpublished report prepared for Nevada Division of Forestry, Carson City, Nevada, and Nevada Fish and Wildlife Office, Reno, Nevada. 48 pp.
- [Service] U.S. Fish and Wildlife Service. 1986. Endangered and threatened wildlife and plants; determination of endangered status for *Eriogonum ovalifolium* var. *williamsiae* (steamboat buckwheat). Federal Register 51: 24669–24672.
- [Service] U.S. Fish and Wildlife Service. 1995. Steamboat buckwheat recovery plan. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon. 33 pp. plus appendices.
- [Service] U.S. Fish and Wildlife Service. 2009. Steamboat buckwheat (*Eriogonum ovalifolium* var. *williamsiae*) 5-year review: summary and evaluation. Nevada Fish and Wildlife Office, Reno, Nevada. 26 pp.
- Schaffer, M.L., and B.A. Stein. 2000. Safeguarding our precious heritage. Pages 301-321 in B.A. Stein, L.S. Kutner, and J.S. Adams (editors), Precious heritage: the status of biodiversity in the United States. Oxford University Press, New York.
- Washoe County. 2018. Washoe County Real Property Assessment Data [web application]. Reno, Nevada. Available online at <  
<https://www.washoecounty.us/assessor/cama/?parid=4945051&Card=1&disclaimer=yes>  
>. Accessed 24 September 2018.



## APPENDIX A: Monitoring Results for *Eriogonum ovalifolium* var. *williamsiae*

Appendix A describes results from 2003, 2005, and 2017 monitoring of *Eriogonum ovalifolium* var. *williamsiae* in the Central Drainage and Main Terrace permanent transects established in 2003. The Central Drainage is a shallow, ephemeral drainage basin with deep colluvial sinter soils that support high densities of *E. ovalifolium* var. *williamsiae*. It contrasts sharply with the Main Terrace, which is a largely un-vegetated expanse of exposed sinter bedrock with little soil development and smaller numbers of *E. ovalifolium* var. *williamsiae* individuals. Until the mid-1980s, the Main Terrace was the location of several geysers that expelled boiling waters dozens of meters into the air. The demographic monitoring program and sampling procedure is described in Pavlik and Stanton 2003. The monitoring program specifies that transects should be monitored at 5 and 10 years post-installation in 2003. However, an interim monitoring event occurred in 2005 to record a possible recruitment event due to a very wet precipitation year. Monitoring did not occur again until 2017 – 14 years post-installation.

### Data Analysis

For the Central Drainage, in 2003, *Eriogonum ovalifolium* var. *williamsiae* data were collected along 16 established transects (0, 6, 10, 17, 25, 27, 35, 37, 44, 49, 51, 53, 87, 89, 94, 99). In 2005, data were collected along 8 of the 16 transects (0, 6, 35, 37, 87, 89, 94, 99). In 2017, data were collected along 16 transects (0, 6, 10, 17, 25, 27, 35, 37, 44, 46, 51, 53, 87, 89, 94, 99); however, one transect differed from what was established in 2003. For the purposes of this analysis, data was analyzed on 7 of the 16 transects that were consistent across all three years. For 2003, 2005, and 2017, weed cover data was collected and analyzed for five transects (17, 35, 46, 51, 87). For 2003 and 2017, vegetative functional group cover data was collected and analyzed along 16 transects listed above for 2017.

For the Main Terrace, in 2003, *Eriogonum ovalifolium* var. *williamsiae* data were collected along 20 established transects (1, 4, 16, 19, 23, 28, 32, 37, 41, 46, 50, 52, 60, 69, 72, 75, 86, 88, 91, 95). In 2005 and 2017, data were collected along these same 20 transects. For the purposes of this analysis, data was analyzed for all 20 transects because they were consistent across all three years. For 2003, 2005, and 2017, weed cover data was collected and analyzed for five transects (16, 32, 50, 69, 86). For 2003 and 2017, vegetative functional group cover data was collected and analyzed for the same 20 transects listed above.

Analyses were conducted using JMP version 14 statistical software (SAS Institute Inc., Cary, NC). For *Eriogonum ovalifolium* var. *williamsiae* data, differences in size class (<1 cm – seedling; 1-5 cm – small; 6-10 cm – medium; and >11 cm – large) were compared using a linear model to determine significance of a year (a fixed factor) by site. For weed (*Bromus tectorum*, *Vulpia myuros*, and other weeds) and vegetative functional group data (Tree/shrub, grass, and *E. ovalifolium* var. *williamsiae*), differences in percent cover were compared using a mixed model and the restricted maximum likelihood method to determine a significance of year (fixed) and transect (random) factors by site and species and/or functional group. All values presented are means and standard errors, and significance is determined by P values <0.05.

## Central Drainage

Across all size classes on the Central Drainage, the number of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* plants has significantly differed over time (Figure 1). The number of *E. ovalifolium* var. *williamsiae* plants in the small size class (1-5 cm;  $F_{2,18} = 4.1$ ,  $P=0.033$ ) and medium size class (6-10 cm;  $F_{2,18} = 5.2$ ,  $P=0.016$ ) has decreased since the baseline survey in 2003. The number of *E. ovalifolium* var. *williamsiae* in the large size class (> 11 cm;  $F_{2,18} = 27.8$ ,  $P<0.001$ ) since the baseline survey in 2003. However, since 2005, the number of small plants (1-5 cm) decreased by 57%, the number of medium plants (6-10 cm) decreased by 43%, and the number of large plants (>11 cm) decreased by 27% (Figure 1). Additionally, 290 seedlings (<1 cm) were observed in 2005, perhaps documenting a rare recruitment event for *E. ovalifolium* var. *williamsiae* due to an increased precipitation event. There were three seedlings observed in 2003 and nine seedlings observed in 2017.




Figure 1. Change in size class distribution of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* in the Central Drainage in 2003, 2005, and 2017. Results are from a linear model using the restricted maximum likelihood method to determine a significance of a fixed factor (year) by size class of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* plants. Values are means ( $n=55$ ) for each size class and standard errors. \* indicates significant ( $P < 0.05$ ) differences among years, by size class.

Weed cover data in the Central Drainage is low, with less than 3% cover of all species. *Bromus tectorum* ( $F_{2,158} = 6.5$ ,  $P=0.0019$ ) cover has significantly increased over time. *Vulpia myuros* ( $F_{2,158} = 3.14$ ,  $P=0.0457$ ) cover has significantly decreased over time. Other weed species only covered <1% of the Central Terrace in 2005 ( $F_{2,158} = 1.0$ ,  $P=0.3702$ ; Figure 2).




Figure 2. Percent cover of weed species in the Central Drainage in 2003, 2005, and 2017. Results are from a mixed model using the restricted maximum likelihood method to determine a significance of a fixed (year) and random (transect) factors by weed species. Values are means (n=55) and standard errors. \* indicates significant ( $P < 0.05$ ) differences among years, by functional group or species.

Tree/shrub ( $F_{1,15} = 0.5644$ ,  $P = 0.4641$ ) and grass ( $F_{1,15} = 0.0079$ ,  $P = 0.9302$ ) cover has not significantly changed over time. The cover of *Eriogonum ovalifolium* var. *williamsiae* has increased over time ( $F_{1,15} = 7.76$ ,  $P = 0.0138$ ; Figure 3).




Figure 3. Percent cover of vegetative functional groups in the Central Drainage in 2003, 2005, and 2017. Results are from a mixed model using the restricted maximum likelihood method to determine a significance of a fixed (year) and random (transect) factors by weed species. Values are means (n=16) and standard errors. \* indicates significant ( $P < 0.05$ ) differences among years, by functional group or species.

## Main Terrace

Only the medium size class (6-10 cm) of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* plants on the Main Terrace has significantly differed over time (Figure 4). Though not significant, the number of *E. ovalifolium* var. *williamsiae* plants in the smallest size class (1-5 cm;  $F_{2,57}= 1.38$ ,  $P=0.2587$ ) has decreased since the baseline survey in 2003. The medium size class (6-10 cm;  $F_{2,57} = 4.57$ ,  $P=0.0143$ ) of *E. ovalifolium* var. *williamsiae* plants has decreased since the baseline survey in 2003. Though not significant, the number of *E. ovalifolium* var. *williamsiae* plants in the largest size class (> 11 cm;  $F_{2,57}= 1.1$ ,  $P=0.3393$ ) has increased since both the baseline survey in 2003 and the 2005. Since 2005, the number of small plants (1-5 cm) decreased by 17%, the number of medium plants (6-10 cm) decreased by 25%, but the number of large plants (>11 cm) increased by 10% (Figure 4). Only one seedling (<1 cm) of *E. ovalifolium* var. *williamsiae* was found during sampling of the Main Terrace in 2003; seedlings were not observed in 2005 and 2017.




Figure 4. Change in size class distribution of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* in the Main Terrace in 2003, 2005, and 2017. Results are from a linear model using the restricted maximum likelihood method to determine a significance of a fixed factor (year) by size class of living (reproductive and non-reproductive) *Eriogonum ovalifolium* var. *williamsiae* plants. Values are means ( $n=20$ ) for each size class and standard errors. \* indicates significant ( $P < 0.05$ ) differences among years, by size class.

Weed cover data in the Main Terrace is low, with less than 3% cover of all species. *Bromus tectorum* (cheatgrass;  $F_{2,158}= 0.3$ ,  $P=0.6915$ ) cover has remained stable over time. The cover of *Vulpia myuros* (rattail fescue;  $F_{2,158}= 4.78$ ,  $P=0.0096$ ) and other weeds ( $F_{2,158}= 32.47$ ,  $P<0.0001$ ) have significantly increased over time (Figure 5).




Figure 5. Percent cover of weed species in the Main Terrace in 2003, 2005, and 2017. Results are from a mixed model using the restricted maximum likelihood method to determine a significance of a fixed (year) and random (transect) factors by weed species. Values are means (n=55) and standard errors. \* indicates significant ( $P < 0.05$ ) differences among years, by functional group or species.

Tree/shrub ( $F_{1,19} = 1.7071$ ,  $P = 0.2070$ ) and grass ( $F_{1,19} = 0.7016$ ,  $P = 0.4126$ ) cover has not significantly changed over time, continuing to remain sparse. The cover of *Eriogonum ovalifolium* var. *williamsiae* has increased, but not significantly, over time ( $F_{1,19} = 0.4997$ ,  $P = 0.4882$ ; Figure 6).




Figure 6. Percent cover of vegetative functional groups in the Main Terrace in 2003, 2005, and 2017. Results are from a mixed model using the restricted maximum likelihood method to determine a significance of a fixed (year) and random (transect) factors by weed species. Values are means (n=20) and standard errors.